

Abstract Submitted
for the OSS16 Meeting of
The American Physical Society

Degree of Dependence and Independence of Neutrosophic Logic Components Applied in Physics FLORENTIN SMARANDACHE, University of New Mexico — Neutrosophic Logic is a general framework for unification of many existing logics, and its components T (truth), I (indeterminacy), F (falsehood) are standard or non-standard real subsets of $]^{-0, 1^{+}[$ with not necessarily any connection between them. For single valued neutrosophic logic, the sum of the components (T+I+F) is: $0 \leq T+I+F \leq 3$ when all three components are independent; $0 \leq T+I+F \leq 2$ when two components are dependent, while the third one is independent from them; $0 \leq T+I+F \leq 1$ when all three components are dependent. When three or two of the components T, I, F are independent, one leaves room for incomplete information (sum <1), paraconsistent and contradictory information (sum >1), or complete information (sum = 1). If all three components T, I, F are dependent, then similarly one leaves room for incomplete information (sum <1), or complete information (sum = 1). The dependent components are tied together. Three sources that provide information on T, I, and F respectively are independent if they do not communicate with each other and do not influence each other. The sum of two components x and y in general is: $0 \leq x+y \leq 2 - d(x, y)$, where $d(x, y)$ is the *degree of dependence* between x and y. Therefore $2 - d(x, y)$ is the *degree of independence* between x and y. But $\max\{T+I+F\}$ may also get any value in $[1, 3]$. For example, suppose that T and F are 30% dependent and 70% independent (hence $T+F \leq 2-0.3 = 1.7$), while I and F are 60% dependent and 40% independent (hence $I+F \leq 2-0.6 = 1.4$). Then $\max\{T+I+F\} = 2.4$ and occurs for $T = 1, I = 0.7, F = 0.7$. p.).

Florentin Smarandache
University of New Mexico

Date submitted: 18 Mar 2016

Electronic form version 1.4