

**ON IMPROVEMENT IN ESTIMATING POPULATION PARAMETER(S) USING
AUXILIARY INFORMATION**

Rajesh Singh

Department of Statistics, BHU, Varanasi (U. P.), India

Editor

Florentin Smarandache

Department of Mathematics, University of New Mexico, Gallup, USA

Editor

<i>Estimator</i>	<i>Bias</i>		<i>MSE</i>	
	<i>First order</i>	<i>Second order</i>	<i>First order</i>	<i>Second order</i>
<i>t₁</i>	0.0044915	0.004424	39.217225	39.45222
<i>t₂</i>	0	-0.00036	39.217225 (for g=1)	39.33552 (for g=1)
<i>t₃</i>	-0.04922	-0.04935	39.217225	39.29102
<i>t₄</i>	0.2809243	-0.60428	39.217225	39.44855
<i>t₅</i>	-0.027679	-0.04911	39.217225	39.27187

**ON IMPROVEMENT IN ESTIMATING POPULATION PARAMETER(S) USING
AUXILIARY INFORMATION**

Rajesh Singh

Department of Statistics, BHU, Varanasi (U. P.), India

Editor

Florentin Smarandache

Department of Mathematics, University of New Mexico, Gallup, USA

Editor

Educational Publishing & Journal of Matter Regularity (Beijing)

2013

This book can be ordered on paper or electronic formats from:

*Education Publishing
1313 Chesapeake Avenue
Columbus, Ohio 43212
USA
Tel. (614) 485-0721
E-mail: info@edupublisher.com*

*Copyright 2013 by EducationPublishing & Journal of Matter Regularity (Beijing), Editors
and the Authors for their papers*

Front and back covers by the Editors

Peer Reviewers:

Prof. Adrian Olaru, University POLITEHNICA of Bucharest (UPB), Bucharest, Romania.

*Dr. Linfan Mao, Academy of Mathematics and Systems, Chinese Academy of Sciences,
Beijing 100190, P. R. China.*

*Y. P. Liu, Department of Applied Mathematics, Beijing Jiaotong University, Beijing 100044,
P. R. China.*

*Prof. Xingsen Li, School of Management, Ningbo Institute of Technology, Zhejiang
University, Ningbo 315100, P. R. China.*

ISBN: 9781599732305

Printed in the United States of America

Contents

Preface: 4

- 1. Use of auxiliary information for estimating population mean in systematic sampling under non- response: 5**
- 2. Some improved estimators of population mean using information on two auxiliary attributes: 17**
- 3. Study of some improved ratio type estimators under second order approximation: 25**
- 4. Improvement in estimating the population mean using dual to ratio-cum-product estimator in simple random sampling: 42**
- 5. Some improved estimators for population variance using two auxiliary variables in double sampling: 54**

Preface

The purpose of writing this book is to suggest some improved estimators using auxiliary information in sampling schemes like simple random sampling and systematic sampling.

This volume is a collection of five papers, written by eight coauthors (listed in the order of the papers): Manoj K. Chaudhary, Sachin Malik, Rajesh Singh, Florentin Smarandache, Hemant Verma, Prayas Sharma, Olufadi Yunusa, and Viplav Kumar Singh, from India, Nigeria, and USA.

The following problems have been discussed in the book:

In chapter one an estimator in systematic sampling using auxiliary information is studied in the presence of non-response. In second chapter some improved estimators are suggested using auxiliary information. In third chapter some improved ratio-type estimators are suggested and their properties are studied under second order of approximation.

In chapter four and five some estimators are proposed for estimating unknown population parameter(s) and their properties are studied.

This book will be helpful for the researchers and students who are working in the field of finite population estimation.

Use of Auxiliary Information for Estimating Population Mean in Systematic Sampling under Non- Response

¹Manoj K. Chaudhary, ¹Sachin Malik, †¹Rajesh Singh and ²Florentin Smarandache

¹Department of Statistics, Banaras Hindu University, Varanasi-221005, India

²Department of Mathematics, University of New Mexico, Gallup, USA

†*Corresponding author*

Abstract

In this paper we have adapted Singh and Shukla (1987) estimator in systematic sampling using auxiliary information in the presence of non-response. The properties of the suggested family have been discussed. Expressions for the bias and mean square error (MSE) of the suggested family have been derived. The comparative study of the optimum estimator of the family with ratio, product, dual to ratio and sample mean estimators in systematic sampling under non-response has also been done. One numerical illustration is carried out to verify the theoretical results.

Keywords: Auxiliary variable, systematic sampling, factor-type estimator, mean square error, non-response.

1. Introduction

There are some natural populations like forests etc., where it is not possible to apply easily the simple random sampling or other sampling schemes for estimating the population characteristics. In such situations, one can easily implement the method of systematic sampling for selecting a sample from the population. In this sampling scheme, only the first unit is selected at random, the rest being automatically selected according to a predetermined pattern. Systematic sampling has been considered in detail by Madow and Madow (1944), Cochran (1946) and Lahiri (1954). The application of systematic sampling to forest surveys has been illustrated by Hasel (1942), Finney (1948) and Nair and Bhargava (1951).

The use of auxiliary information has been permeated the important role to improve the efficiency of the estimators in systematic sampling. Kushwaha and Singh (1989) suggested a class of almost unbiased ratio and product type estimators for estimating the population mean using jack-knife technique initiated by Quenouille (1956). Later Banarasi et al. (1993), Singh and Singh (1998), Singh et al. (2012), Singh et al. (2012) and Singh and Solanki (2012) have made an attempt to improve the estimators of population mean using auxiliary information in systematic sampling.

The problem of non-response is very common in surveys and consequently the estimators may produce bias results. Hansen and Hurwitz (1946) considered the problem of estimation of population mean under non-response. They proposed a sampling plan that involves taking a subsample of non-respondents after the first mail attempt and then enumerating the subsample by personal interview. El-Badry (1956) extended Hansen and Hurwitz (1946) technique. Hansen and Hurwitz (1946) technique in simple random sampling is described as: From a population $U = (U_1, U_2, \dots, U_N)$, a large first phase sample of size n' is selected by simple random sampling without replacement (SRSWOR). A smaller second phase sample of size n is selected from n' by SRSWOR. Non-response occurs on the second phase of size n in which n_1 units respond and n_2 units do not. From the n_2 non-respondents, by SRSWOR a sample of $r = n_2/k$; $k > 1$ units is selected. It is assumed that all the r units respond this time round. (see Singh and Kumar (20009)). Several authors such as Cochran (1977), Sodipo and Obisesan (2007), Rao (1987), Khare and Srivastava (1997) and Okafor and Lee (2000) have studied the problem of non-response under SRS.

In the sequence of improving the estimator, Singh and Shukla (1987) proposed a family of factor-type estimators for estimating the population mean in simple random sampling using an auxiliary variable, as

$$T_{\alpha} = \bar{y} \left[\frac{(A+C)\bar{X} + fB\bar{x}}{(A+fB)\bar{X} + C\bar{x}} \right] \quad (1.1)$$

where \bar{y} and \bar{x} are the sample means of the population means \bar{Y} and \bar{X} respectively. A , B and C are the functions of α , which is a scalar and chosen so as the MSE of the estimator T_{α} is minimum.

Where,

$$A = (\alpha - 1)(\alpha - 2), \quad B = (\alpha - 1)(\alpha - 4),$$

$$C = (\alpha - 2)(\alpha - 3)(\alpha - 4); \quad \alpha > 0 \text{ and}$$

$$f = \frac{n}{N}.$$

Remark 1 : If we take $\alpha = 1, 2, 3$ and 4 , the resulting estimators will be ratio, product, dual to ratio and sample mean estimators of population mean in simple random sampling respectively (for details see Singh and Shukla (1987)).

In this paper, we have proposed a family of factor-type estimators for estimating the population mean in systematic sampling in the presence of non-response adapting Singh and Shukla (1987) estimator. The properties of the proposed family have been discussed with the help of empirical study.

2. Sampling Strategy and Estimation Procedure

Let us assume that a population consists of N units numbered from 1 to N in some order. If $N = nk$, where k is a positive integer, then there will be k possible samples each consisting of n units. We select a sample at random and collect the information from the units of the selected sample. Let n_1 units in the sample responded and n_2 units did not respond, so that $n_1 + n_2 = n$. The n_1 units may be regarded as a sample from the response class and n_2 units as a sample from the non-response class belonging to the population. Let us assume that N_1 and N_2 be the number of units in the response class and non-response

class respectively in the population. Obviously, N_1 and N_2 are not known but their unbiased estimates can be obtained from the sample as

$$\hat{N}_1 = n_1 N / n; \quad \hat{N}_2 = n_2 N / n.$$

Further, using Hansen and Hurwitz (1946) technique we select a sub-sample of size h_2 from the n_2 non-respondent units such that $n_2 = h_2 L$ ($L > 1$) and gather the information on all the units selected in the sub-sample (for details on Hansen and Hurwitz (1946) technique see Singh and Kumar (2009)).

Let Y and X be the study and auxiliary variables with respective population means \bar{Y} and \bar{X} . Let $y_{ij}(x_{ij})$ be the observation on the j^{th} unit in the i^{th} systematic sample under study (auxiliary) variable ($i = 1 \dots k : j = 1 \dots n$). Let us consider the situation in which non-response is observed on study variable and auxiliary variable is free from non-response. The Hansen-Hurwitz (1946) estimator of population mean \bar{Y} and sample mean estimator of \bar{X} based on a systematic sample of size n , are respectively given by

$$\bar{y}^* = \frac{n_1 \bar{y}_{n_1} + n_2 \bar{y}_{h_2}}{n}$$

$$\text{and } \bar{x} = \frac{1}{n} \sum_{j=1}^n x_{ij}$$

where \bar{y}_{n_1} and \bar{y}_{h_2} are respectively the means based on n_1 respondent units and h_2 non-respondent units. Obviously, \bar{y}^* and \bar{x} are unbiased estimators of \bar{Y} and \bar{X} respectively. The respective variances of \bar{y}^* and \bar{x} are expressed as

$$V(\bar{y}^*) = \frac{N-1}{nN} \{1 + (n-1)\rho_Y\} S_Y^2 + \frac{L-1}{n} W_2 S_{Y_2}^2 \quad (2.1)$$

and

$$V(\bar{x}) = \frac{N-1}{nN} \{1 + (n-1)\rho_X\} S_X^2 \quad (2.2)$$

where ρ_Y and ρ_X are the correlation coefficients between a pair of units within the systematic sample for the study and auxiliary variables respectively. S_Y^2 and S_X^2 are respectively the mean squares of the entire group for study and auxiliary variables. $S_{Y_2}^2$ be the population mean square of non-response group under study variable and W_2 is the non-response rate in the population.

Assuming population mean \bar{X} of auxiliary variable is known, the usual ratio, product and dual to ratio estimators based on a systematic sample under non-response are respectively given by

$$\bar{y}_R^* = \frac{\bar{y}}{\bar{x}} \bar{X}, \quad (2.3)$$

$$\bar{y}_P^* = \frac{\bar{y} \bar{x}}{\bar{X}} \quad (2.4)$$

and
$$\bar{y}_D^* = \bar{y} \frac{(N\bar{X} - n\bar{x})}{(N-n)\bar{X}}. \quad (2.5)$$

Obviously, all the above estimators \bar{y}_R^* , \bar{y}_P^* and \bar{y}_D^* are biased. To derive the biases and mean square errors (MSE) of the estimators \bar{y}_R^* , \bar{y}_P^* and \bar{y}_D^* under large sample approximation, let

$$\bar{y}^* = \bar{Y}(1 + e_0)$$

$$\bar{x} = \bar{X}(1 + e_1)$$

such that $E(e_0) = E(e_1) = 0$,

$$E(e_0^2) = \frac{V(\bar{y}^*)}{\bar{Y}^2} = \frac{N-1}{nN} \{1 + (n-1)\rho_Y\} C_Y^2 + \frac{L-1}{n} W_2 \frac{S_{Y_2}^2}{\bar{Y}^2}, \quad (2.6)$$

$$E(e_1^2) = \frac{V(\bar{x})}{\bar{X}^2} = \frac{N-1}{nN} \{1 + (n-1)\rho_X\} C_X^2 \quad (2.7)$$

and

$$E(e_0e_1) = \frac{Cov(\bar{y}^*, \bar{x})}{\bar{Y}\bar{X}} = \frac{N-1}{nN} \{1+(n-1)\rho_Y\}^{1/2} \{1+(n-1)\rho_X\}^{1/2} \rho C_Y C_X \quad (2.8)$$

where C_Y and C_X are the coefficients of variation of study and auxiliary variables respectively in the population (for proof see Singh and Singh(1998) and Singh (2003, pg. no. 138)).

The biases and MSE's of the estimators \bar{y}_R^* , \bar{y}_P^* and \bar{y}_D^* up to the first order of approximation using (2.6-2.8), are respectively given by

$$B(\bar{y}_R^*) = \frac{N-1}{nN} \bar{Y} \{1+(n-1)\rho_X\} (1-K\rho^*) C_X^2, \quad (2.9)$$

$$MSE(\bar{y}_R^*) = \frac{N-1}{nN} \bar{Y}^2 \{1+(n-1)\rho_X\} [\rho^{*2} C_Y^2 + (1-2K\rho^*) C_X^2] + \frac{L-1}{n} W_2 S_{Y2}^2, \quad (2.10)$$

$$B(\bar{y}_P^*) = \frac{N-1}{nN} \bar{Y} \{1+(n-1)\rho_X\} K\rho^* C_X^2, \quad (2.11)$$

$$MSE(\bar{y}_P^*) = \frac{N-1}{nN} \bar{Y}^2 \{1+(n-1)\rho_X\} [\rho^{*2} C_Y^2 + (1+2K\rho^*) C_X^2] + \frac{L-1}{n} W_2 S_{Y2}^2, \quad (2.12)$$

$$B(\bar{y}_D^*) = \frac{N-1}{nN} \bar{Y} \{1+(n-1)\rho_X\} [-\rho^* K] C_X^2, \quad (2.13)$$

$$MSE(\bar{y}_D^*) = \frac{N-1}{nN} \bar{Y}^2 \{1+(n-1)\rho_X\} \left[\rho^{*2} C_Y^2 + \left(\frac{f}{1-f} \right) \left\{ \left(\frac{f}{1-f} \right) - 2\rho^* K \right\} C_X^2 \right] + \frac{(L-1)}{n} W_2 S_{Y2}^2 \quad (2.14)$$

where,

$$\rho^* = \frac{\{1+(n-1)\rho_Y\}^{1/2}}{\{1+(n-1)\rho_X\}^{1/2}} \quad \text{and} \quad K = \rho \frac{C_Y}{C_X}.$$

(for details of proof refer to Singh et al.(2012)).

The regression estimator based on a systematic sample under non-response is given by

$$\bar{y}_{lr}^* = \bar{y}^* + b(\bar{X} - \bar{x}) \quad (2.15)$$

MSE of the estimator \bar{y}_{lr}^* is given by

$$\text{MSE}(\bar{y}_{lr}^*) = \frac{N-1}{nN} \bar{Y}^2 \{1 + (n-1)\rho_x\} [C_y^2 - K^2 C_x^2] \rho^{*2} + \frac{(L-1)}{n} W_2 S_{Y2}^2 \quad (2.16)$$

3. Adapted Family of Estimators

Adapting the estimator proposed by Singh and Shukla (1987), a family of factor-type estimators of population mean in systematic sampling under non-response is written as

$$T_\alpha^* = \bar{y}^* \left[\frac{(A+C)\bar{X} + fB\bar{x}}{(A+fB)\bar{X} + C\bar{x}} \right]. \quad (3.1)$$

The constants A, B, C, and f are same as defined in (1.1).

It can easily be seen that the proposed family generates the non-response versions of some well known estimators of population mean in systematic sampling on putting different choices of α . For example, if we take $\alpha = 1, 2, 3$ and 4 , the resulting estimators will be ratio, product, dual to ratio and sample mean estimators of population mean in systematic sampling under non-response respectively.

3.1 Properties of T_α^*

Obviously, the proposed family is biased for the population mean \bar{Y} . In order to find the bias and MSE of T_α^* , we use large sample approximations. Expressing the equation (3.1) in terms of e_i 's ($i = 0,1$) we have

$$T_\alpha^* = \frac{\bar{Y}(1+e_0)(1+De_1)^{-1}[(A+C)+fB(1+e_1)]}{A+fB+C} \quad (3.2)$$

$$\text{where } D = \frac{C}{A+fB+C}.$$

Since $|D| < 1$ and $|e_i| < 1$, neglecting the terms of e_i 's ($i = 0,1$) having power greater than two, the equation (3.2) can be written as

$$T_{\alpha}^* - \bar{Y} = \frac{\bar{Y}}{A + fB + C} \left[(A + C) \{e_0 - De_1 + D^2e_1^2 - De_0e_1\} + fB \{e_0 - (D-1)e_1 + D(D-1)e_1^2 - (D-1)e_0e_1\} \right]. \quad (3.3)$$

Taking expectation of both sides of the equation (3.3), we get

$$E[T_{\alpha}^* - \bar{Y}] = \frac{\bar{Y}(C - fB)}{A + fB + C} \left[\frac{C}{A + fB + C} E(e_1^2) - E(e_0e_1) \right].$$

Let $\phi_1(\alpha) = \frac{fB}{A + fB + C}$ and $\phi_2(\alpha) = \frac{C}{A + fB + C}$ then

$$\phi(\alpha) = \phi_2(\alpha) - \phi_1(\alpha) = \frac{C - fB}{A + fB + C}.$$

Thus, we have

$$E[T_{\alpha}^* - \bar{Y}] = \bar{Y}\phi(\alpha) [\phi_2(\alpha)E(e_1^2) - E(e_0e_1)]. \quad (3.4)$$

Putting the values of $E(e_1^2)$ and $E(e_0e_1)$ from equations (2.7) and (2.8) into the equation (3.4), we get the bias of T_{α}^* as

$$B(T_{\alpha}^*) = \phi(\alpha) \frac{N-1}{nN} \bar{Y} \{1 + (n-1)\rho_x\} [\phi_2(\alpha) - \rho^* K] C_x^2. \quad (3.5)$$

Squaring both the sides of the equation (3.3) and then taking expectation, we get

$$E[T_{\alpha}^* - \bar{Y}]^2 = \bar{Y}^2 [E(e_0^2) + \phi^2(\alpha)E(e_1^2) - 2\phi(\alpha)E(e_0e_1)]. \quad (3.6)$$

Substituting the values of $E(e_0^2)$, $E(e_1^2)$ and $E(e_0e_1)$ from the respective equations (2.6), (2.7) and (2.8) into the equation (3.6), we get the MSE of T_{α}^* as

$$MSE(T_{\alpha}^*) = \frac{N-1}{nN} \bar{Y}^2 \{1 + (n-1)\rho_x\} \left[\rho^{*2} C_Y^2 + \{\phi^2(\alpha) - 2\phi(\alpha)\rho^* K\} C_x^2 \right] + \frac{(L-1)}{n} W_2 S_{Y2}^2. \quad (3.7)$$

3.2 Optimum Choice of α

In order to obtain the optimum choice of α , we differentiate the equation (3.7) with respect to α and equating the derivative to zero, we get the normal equation as

$$\frac{N-1}{nN} \bar{Y}^2 \{1 + (n-1)\rho_x\} [2\phi(\alpha)\phi'(\alpha) - 2\phi'(\alpha)\rho^*K] C_x^2 = 0 \quad (3.8)$$

where $\phi'(\alpha)$ is the first derivative of $\phi(\alpha)$ with respect to α .

Now from equation (3.8), we get

$$\phi(\alpha) = \rho^*K \quad (3.9)$$

which is the cubic equation in α . Thus α has three real roots for which the MSE of proposed family would attain its minimum.

Putting the value of $\phi(\alpha)$ from equation (3.9) into equation (3.7), we get

$$MSE(T_{\alpha}^*)_{\min} = \frac{N-1}{nN} \bar{Y}^2 \{1 + (n-1)\rho_x\} [C_Y^2 - K^2 C_X^2] \rho^{*2} + \frac{(L-1)}{n} W_2 S_{Y2}^2 \quad (3.10)$$

which is the MSE of the usual regression estimator of population mean in systematic sampling under non-response.

4. Empirical Study

In the support of theoretical results, we have considered the data given in Murthy (1967, p. 131-132). These data are related to the length and timber volume for ten blocks of the blacks mountain experimental forest. The value of intraclass correlation coefficients ρ_x and ρ_y have been given approximately equal by Murthy (1967, p. 149) and Kushwaha and Singh (1989) for the systematic sample of size 16 by enumerating all possible systematic samples after arranging the data in ascending order of strip length. The particulars of the population are given below:

$$\begin{aligned} N = 176, \quad n = 16, \quad \bar{Y} = 282.6136, \quad \bar{X} = 6.9943, \\ S_Y^2 = 24114.6700, \quad S_X^2 = 8.7600, \quad \rho = 0.8710, \end{aligned}$$

$$S_{Y_2}^2 = \frac{3}{4} S_Y^2 = 18086.0025.$$

Table 1 depicts the MSE's and variance of the estimators of proposed family with respect to non-response rate (W_2).

Table 1: MSE and Variance of the Estimators for $L = 2$.

α	W_2			
	0.1	0.2	0.3	0.4
1 ($= \bar{y}_R^*$)	371.37	484.41	597.45	710.48
2 ($= \bar{y}_P^*$)	1908.81	2021.85	2134.89	2247.93
3 ($= \bar{y}_D^*$)	1063.22	1176.26	1289.30	1402.33
4 ($= \bar{y}^*$)	1140.69	1253.13	1366.17	1479.205
$\alpha_{opt} (= (T_\alpha^*)_{min})$	270.67	383.71	496.75	609.78

5. Conclusion

In this paper, we have adapted Singh and Shukla (1987) estimator in systematic sampling in the presence of non-response using an auxiliary variable and obtained the optimum estimator of the proposed family. It is observed that the proposed family can generate the non-response versions of a number of estimators of population mean in systematic sampling on different choice of α . From Table 1, we observe that the proposed family under optimum condition has minimum MSE, which is equal to the MSE of the regression estimator (most of the class of estimators in sampling literature under optimum condition attains MSE equal to the MSE of the regression estimator). It is also seen that the MSE or variance of the estimators increases with increase in non response rate in the population.

References

1. Banarasi, Kushwaha, S.N.S. and Kushwaha, K.S. (1993): A class of ratio, product and difference (RPD) estimators in systematic sampling, *Microelectron. Reliab.*, 33, 4, 455–457.
2. Cochran, W. G. (1946): Relative accuracy of systematic and stratified random samples for a certain class of population, *AMS*, 17, 164-177.
3. Finney, D.J. (1948): Random and systematic sampling in timber surveys, *Forestry*, 22, 64-99.
4. Hansen, M. H. and Hurwitz, W. N. (1946) : The problem of non-response in sample surveys, *Jour. of The Amer. Stat. Assoc.*, 41, 517-529.
5. Hasel, A. A. (1942): Estimation of volume in timber stands by strip sampling, *AMS*, 13, 179-206.
6. Kushwaha, K. S. and Singh, H.P. (1989): Class of almost unbiased ratio and product estimators in systematic sampling, *Jour. Ind. Soc. Ag. Statistics*, 41, 2, 193–205.
7. Lahiri, D. B. (1954): On the question of bias of systematic sampling, *Proceedings of World Population Conference*, 6, 349-362.
8. Madow, W. G. and Madow, L.H. (1944): On the theory of systematic sampling, *I. Ann. Math. Statist.*, 15, 1-24.
9. Murthy, M.N. (1967): *Sampling Theory and Methods*. Statistical Publishing Society, Calcutta.
10. Nair, K. R. and Bhargava, R. P. (1951): Statistical sampling in timber surveys in India, *Forest Research Institute, Dehradun, Indian forest leaflet*, 153.
11. Quenouille, M. H. (1956): Notes on bias in estimation, *Biometrika*, 43, 353-360.
12. Singh, R and Singh, H. P. (1998): Almost unbiased ratio and product type- estimators in systematic sampling, *Questiio*, 22,3, 403-416.
13. Singh, R., Malik, S., Chaudhary, M.K., Verma, H. and Adewara, A. A. (2012) : A general family of ratio type- estimators in systematic sampling. *Jour. Reliab. and Stat. Stud.*,5(1), 73-82).
14. Singh, R., Malik, S., Singh, V. K. (2012) : An improved estimator in systematic sampling. *Jour. Of Scie. Res.*, 56, 177-182.

15. Singh, H.P. and Kumar, S. (2009) : A general class of dss estimators of population ratio, product and mean in the presence of non-response based on the sub-sampling of the non-respondents. *Pak J. Statist.*, 26(1), 203-238.
16. Singh, H.P. and Solanki, R. S. (2012) : An efficient class of estimators for the population mean using auxiliary information in systematic sampling. *Jour. of Stat. Ther. and Pract.*, 6(2), 274-285.
17. Singh, S. (2003) : Advanced sampling theory with applications. Kluwer Academic Publishers.
18. Singh, V. K. and Shukla, D. (1987): One parameter family of factor-type ratio estimators, *Metron*, 45 (1-2), 273-283.

Some Improved Estimators of Population Mean Using Information on Two Auxiliary Attributes

¹Hemant Verma, †¹Rajesh Singh and ²Florentin Smarandache

¹Department of Statistics, Banaras Hindu University, Varanasi-221005, India

²Department of Mathematics, University of New Mexico, Gallup, USA

†Corresponding author

Abstract

In this paper, we have studied the problem of estimating the finite population mean when information on two auxiliary attributes are available. Some improved estimators in simple random sampling without replacement have been suggested and their properties are studied. The expressions of mean squared error's (MSE's) up to the first order of approximation are derived. An empirical study is carried out to judge the best estimator out of the suggested estimators.

Key words: Simple random sampling, auxiliary attribute, point bi-serial correlation, phi correlation, efficiency.

Introduction

The role of auxiliary information in survey sampling is to increase the precision of estimators when study variable is highly correlated with auxiliary variable. But when we talk about qualitative phenomena of any object then we use auxiliary attributes instead of auxiliary variable. For example, if we talk about height of a person then sex will be a good auxiliary attribute and similarly if we talk about particular breed of cow then in this case milk produced by them will be good auxiliary variable.

Most of the times, we see that instead of one auxiliary variable we have information on two auxiliary variables e.g.; to estimate the hourly wages we can use the information on marital status and region of residence (see Gujrati and Sangeetha (2007), page-311).

In this paper, we assume that both auxiliary attributes have significant point bi-serial correlation with the study variable and there is significant phi-correlation (see Yule (1912)) between the auxiliary attributes.

Consider a sample of size n drawn by simple random sampling without replacement (SRSWOR) from a population of size N . let $y_j, \phi_{ij} (i=1,2)$ denote the observations on variable y and $\phi_i (i=1,2)$ respectively for the j^{th} unit ($i=1,2,3,\dots,N$). We note that $\phi_{ij}=1$, if j^{th} unit possesses attribute $\phi_{ij}=0$ otherwise. Let $A_i = \sum_{j=1}^N \phi_{ij}$, $a_i = \sum_{j=1}^n \phi_{ij}$; $i=1,2$ denotes the total number of units in the population and sample respectively, possessing attribute ϕ . Similarly, let $P_i = \frac{A_i}{N}$ and $p_i = \frac{a_i}{n}$; ($i=1,2$) denotes the proportion of units in the population and sample respectively possessing attribute $\phi_i (i=1,2)$.

In order to have an estimate of the study variable y , assuming the knowledge of the population proportion P , Naik and Gupta (1996) and Singh et al. (2007) respectively proposed following estimators:

$$t_1 = \bar{y} \left(\frac{P_1}{p_1} \right) \quad (1.1)$$

$$t_2 = \bar{y} \left(\frac{p_2}{P_2} \right) \quad (1.2)$$

$$t_3 = \bar{y} \exp \left(\frac{P_1 - p_1}{P_1 + p_1} \right) \quad (1.3)$$

$$t_4 = \bar{y} \exp \left(\frac{p_2 - P_2}{p_2 + P_2} \right) \quad (1.4)$$

The bias and MSE expression's of the estimator's t_i ($i=1, 2, 3, 4$) up to the first order of approximation are, respectively, given by

$$B(t_1) = \bar{Y} f_1 C_{p_1}^2 [1 - K_{pb_1}] \quad (1.5)$$

$$B(t_2) = \bar{Y} f_1 K_{pb_2} C_{p_2}^2 \quad (1.6)$$

$$B(t_3) = \bar{Y}f_1 \frac{C_{p_1}^2}{2} \left[\frac{1}{4} - K_{pb_1} \right] \quad (1.7)$$

$$B(t_4) = \bar{Y}f_1 \frac{C_{p_2}^2}{2} \left[\frac{1}{4} + K_{pb_2} \right] \quad (1.8)$$

$$MSE(t_1) = \bar{Y}^2 f_1 [C_y^2 + C_{p_1}^2 (1 - 2K_{pb_1})] \quad (1.9)$$

$$MSE(t_2) = \bar{Y}^2 f_1 [C_y^2 + C_{p_1}^2 (1 + 2K_{pb_2})] \quad (1.10)$$

$$MSE(t_3) = \bar{Y}^2 f_1 \left[C_y^2 + C_{p_1}^2 \left(\frac{1}{4} - K_{pb_1} \right) \right] \quad (1.11)$$

$$MSE(t_4) = \bar{Y}^2 f_1 \left[C_y^2 + C_{p_2}^2 \left(\frac{1}{4} + K_{pb_2} \right) \right] \quad (1.12)$$

where, $f_1 = \frac{1}{n} - \frac{1}{N}$, $S_{\phi_j}^2 = \frac{1}{N-1} \sum_{i=1}^N (\phi_{ji} - P_j)^2$, $S_{y\phi_j} = \frac{1}{N-1} \sum_{i=1}^N (y_i - \bar{Y})(\phi_{ji} - P_j)$,

$$\rho_{pb_j} = \frac{S_{y\phi_j}}{S_y S_{\phi_j}}, \quad C_y = \frac{S_y}{\bar{Y}}, \quad C_{p_j} = \frac{S_{\phi_j}}{P_j}; \quad (j=1,2), \quad K_{pb_1} = \rho_{pb_1} \frac{C_y}{C_{p_1}}, \quad K_{pb_2} = \rho_{pb_2} \frac{C_y}{C_{p_2}}.$$

$s_{\phi_1\phi_2} = \frac{1}{n-1} \sum_{i=1}^n (\phi_{1i} - p_1)(\phi_{2i} - p_2)$ and $\rho_\phi = \frac{S_{\phi_1\phi_2}}{S_{\phi_1} S_{\phi_2}}$ be the sample phi-covariance and phi-

correlation between ϕ_1 and ϕ_2 respectively, corresponding to the population phi-covariance

and phi-correlation $S_{\phi_1\phi_2} = \frac{1}{N-1} \sum_{i=1}^N (\phi_{1i} - P_1)(\phi_{2i} - P_2)$ and $\rho_\phi = \frac{S_{\phi_1\phi_2}}{S_{\phi_1} S_{\phi_2}}$.

In this paper we have proposed some improved estimators of population mean using information on two auxiliary attributes in simple random sampling without replacement. A comparative study is also carried out to compare the optimum estimators with respect to usual mean estimator with the help of numerical data.

2. Proposed Estimators

Following Olkin (1958), we propose an estimator t_5 as

$$t_5 = \bar{y} \left[w_1 \frac{P_1}{p_1} + w_2 \frac{P_2}{p_2} \right] \quad (2.1)$$

where w_1 and w_2 are constants, such that $w_1 + w_2 = 1$.

Consider another estimator t_6 as

$$t_6 = \left[K_{61} \bar{y} + K_{62} (P_1 - p_1) \right] \exp \left[\frac{P_2 - p_2}{P_2 + p_2} \right] \quad (2.2)$$

where K_{61} and K_{62} are constants.

Following Shaoo et al. (1993), we propose another estimator t_7 as

$$t_7 = \bar{y} + K_{71} (P_1 - p_1) + K_{72} (P_2 - p_2) \quad (2.3)$$

where K_{71} and K_{72} are constants.

Bias and MSE of estimators t_5 , t_6 and t_7 :

To obtain the bias and MSE expressions of the estimators t_i ($i = 5, 6, 7$) to the first degree of approximation, we define

$$e_0 = \frac{\bar{y} - \bar{Y}}{\bar{Y}}, \quad e_1 = \frac{p_1 - P_1}{P_1}, \quad e_2 = \frac{p_2 - P_2}{P_2}$$

such that, $E(e_i) = 0$; $i = 0, 1, 2$.

Also,

$$E(e_0^2) = f_1 C_y^2, \quad E(e_1^2) = f_1 C_{p_1}^2, \quad E(e_2^2) = f_1 C_{p_2}^2,$$

$$E(e_0 e_1) = f_1 K_{pb_1} C_{p_1}^2, \quad E(e_0 e_2) = f_1 K_{pb_2} C_{p_2}^2, \quad E(e_1 e_2) = f_1 K_\phi C_{p_2}^2,$$

$$K_{pb_1} = \rho_{pb_1} \frac{C_y}{C_{p_1}}, \quad K_{pb_2} = \rho_{pb_2} \frac{C_y}{C_{p_2}}, \quad K_\phi = \rho_\phi \frac{C_{p_1}}{C_{p_2}}$$

Expressing (2.1) in terms of e 's we have,

$$t_5 = \bar{Y} (1 + e_0) \left[w_1 \frac{P_1}{P_1 (1 + e_1)} + w_2 \frac{P_2}{P_2 (1 + e_2)} \right]$$

$$t_5 = \bar{Y}(1 + e_0) \left[w_1(1 + e_1)^{-1} + w_2(1 + e_2)^{-1} \right] \quad (3.1)$$

Expanding the right hand side of (3.1) and retaining terms up to second degrees of e 's, we have,

$$t_5 = \bar{Y} \left[1 + e_0 - w_1 e_1 - w_2 e_2 + w_1 e_1^2 + w_2 e_2^2 - w_1 e_0 e_1 - w_2 e_0 e_2 \right] \quad (3.2)$$

Taking expectations of both sides of (3.1) and then subtracting \bar{Y} from both sides, we get the bias of estimator t_5 upto the first order of approximation as

$$\text{Bias}(t_5) = \bar{Y} f_1 \left[w_1 C_{p_1}^2 (1 - K_{pb_1}) + w_2 C_{p_2}^2 (1 - K_{pb_2}) \right] \quad (3.3)$$

From (3.2), we have,

$$(t_5 - \bar{Y}) \cong \bar{Y} [e_0 - w_1 e_1 - w_2 e_2] \quad (3.4)$$

Squaring both sides of (3.4) and then taking expectations, we get the MSE of t_5 up to the first order of approximation as

$$\text{MSE}(t_5) = \bar{Y}^2 f_1 \left[C_y^2 + w_1^2 C_{p_1}^2 + w_2^2 C_{p_2}^2 - 2w_1 K_{pb_1} C_{p_1}^2 - 2w_2 K_{pb_2} C_{p_2}^2 + 2w_1 w_2 K_{\phi} C_{p_2}^2 \right] \quad (3.5)$$

Minimization of (3.5) with respect to w_1 and w_2 , we get the optimum values of w_1 and w_2 , as

$$w_{1(\text{opt})} = \frac{K_{pb_1} C_{p_1}^2 - K_{\phi} C_{p_2}^2}{C_{p_1}^2 - K_{\phi} C_{p_2}^2} = w_1^* (\text{say})$$

$$\begin{aligned} w_{2(\text{opt})} &= 1 - w_{1(\text{opt})} \\ &= 1 - \frac{K_{pb_1} C_{p_1}^2 - K_{\phi} C_{p_2}^2}{C_{p_1}^2 - K_{\phi} C_{p_2}^2} \\ &= \frac{C_{p_1}^2 [1 - K_{pb_1}]}{C_{p_1}^2 - K_{\phi} C_{p_2}^2} = w_2^* (\text{say}) \end{aligned}$$

Similarly, we get the bias and MSE expressions of estimator t_6 and t_7 respectively, as

$$\text{Bias}(t_6) = K_{61} \bar{Y} \left[1 + f_1 C_{p_2}^2 \left(\frac{3}{8} - \frac{1}{2} K_{pb_2} \right) \right] + \frac{1}{2} K_{22} P_1 f_1 K_{\phi} C_{p_2}^2 \quad (3.6)$$

$$\text{Bias}(t_7) = 0 \quad (3.7)$$

And

$$\text{MSE}(t_6) = K_{61}^2 \bar{Y}^2 A_1 + K_{62}^2 P_1^2 A_2 - 2K_{61}K_{62}P_1 \bar{Y}A_3 + (1 - 2K_{61})\bar{Y}^2 \quad (3.8)$$

$$\text{where } A_1 = 1 + f_1 \left(C_y^2 + C_{p_2}^2 \left(\frac{1}{4} - K_{pb_2} \right) \right)$$

$$A_2 = f_1 C_{p_1}^2$$

$$A_3 = f_1 \left(k_{pb_1} C_{p_1}^2 - \frac{1}{2} K_{\phi} C_p^2 \right)$$

And the optimum values of K_{61} and K_{62} are respectively, given as

$$K_{61(\text{opt})} = \frac{A_2}{A_1 A_2 - A_3^2} = K_{61}^* (\text{say})$$

$$K_{62(\text{opt})} = \frac{\bar{Y} A_3}{P_1 (A_1 A_2 - A_3^2)} = K_{62}^* (\text{say})$$

$$\begin{aligned} \text{MSE}(t_7) = & \bar{Y}^2 f_1 C_y^2 + K_{71}^2 P_1^2 f_1 C_{p_1}^2 + K_{72}^2 P_2^2 f_1 C_{p_2}^2 - 2K_{71} P_1 \bar{Y} f_1 K_{pb_1} C_{p_1}^2 - 2K_{72} P_2 \bar{Y} f_1 K_{pb_2} C_{p_2}^2 \\ & + 2K_{71} K_{72} P_1 P_2 f_1 K_{\phi} C_p^2 \end{aligned} \quad (3.9)$$

And the optimum values of K_{71} and K_{72} are respectively, given as

$$K_{71(\text{opt})} = \frac{\bar{Y}}{P_1} \left(\frac{K_{pb_1} C_{p_1}^2 - K_{pb_2} K_{\phi} C_{p_2}^2}{C_{p_1}^2 - K_{\phi}^2 C_{p_2}^2} \right) = K_{71}^* (\text{say})$$

$$K_{72(\text{opt})} = \frac{\bar{Y}}{P_2} \left(\frac{K_{pb_2} C_{p_1}^2 - K_{pb_1} K_{\phi} C_{p_1}^2}{C_{p_1}^2 - K_{\phi}^2 C_{p_2}^2} \right) = K_{72}^* (\text{say})$$

4. Empirical Study

Data: (Source: Government of Pakistan (2004))

The population consists rice cultivation areas in 73 districts of Pakistan. The variables are defined as:

Y = rice production (in 000' tonnes, with one tonne = 0.984 ton) during 2003,

P_1 = production of farms where rice production is more than 20 tonnes during the year 2002, and

P_2 = proportion of farms with rice cultivation area more than 20 hectares during the year 2003.

For this data, we have

$$N=73, \bar{Y}=61.3, P_1=0.4247, P_2=0.3425, S_y^2=12371.4, S_{\phi_1}^2=0.225490, S_{\phi_2}^2=0.228311,$$

$$\rho_{pb_1}=0.621, \rho_{pb_2}=0.673, \rho_{\phi}=0.889.$$

The percent relative efficiency (PRE's) of the estimators t_i ($i=1,2,\dots,7$) with respect to unusual unbiased estimator \bar{y} have been computed and given in Table 4.1.

Table 4.1 : PRE of the estimators with respect to \bar{y}

Estimator	PRE
\bar{y}	100.00
t_1	162.7652
t_2	48.7874
t_3	131.5899
t_4	60.2812
t_5	165.8780
t_6	197.7008
t_7	183.2372

Conclusion

In this paper we have proposed some improved estimators of population mean using information on two auxiliary attributes in simple random sampling without replacement. From the Table 4.1 we observe that the estimator t_6 is the best followed by the estimator t_7 .

References

Government of Pakistan, 2004, Crops Area Production by Districts (Ministry of Food, Agriculture and Livestock Division, Economic Wing, Pakistan).

Gujarati, D. N. and Sangeetha (2007): Basic econometrics. Tata McGraw – Hill.

- Malik, S. and Singh, R. (2013): A family of estimators of population mean using information on point bi-serial and phi correlation coefficient. IJSE
- Naik, V. D and Gupta, P.C.(1996): A note on estimation of mean with known population proportion of an auxiliary character. Jour. Ind. Soc. Agri. Stat., 48(2), 151-158.
- Olkin, I. (1958): Multivariate ratio estimation for finite populations, Biometrika, 45, 154–165.
- Singh, R., Chauhan, P., Sawan, N. and Smarandache, F.(2007): Auxiliary information and a priori values in construction of improved estimators. Renaissance High press.
- Singh, R., Chauhan, P., Sawan, N. and Smarandache, F. (2008): Ratio Estimators in Simple Random Sampling Using Information on Auxiliary Attribute. Pak. Jour. Stat. Oper. Res. Vol. IV, No.1, pp. 47-53.
- Singh, R., Kumar, M. and Smarandache, F. (2010): Ratio estimators in simple random sampling when study variable is an attribute. WASJ 11(5): 586-589.
- Yule, G. U. (1912): On the methods of measuring association between two attributes. Jour. of the Royal Soc. 75, 579-642.

Study of Some Improved Ratio Type Estimators Under Second Order Approximation

¹Prayas Sharma, †¹Rajesh Singh and ²Florentin Smarandache

¹Department of Statistics, Banaras Hindu University, Varanasi-221005, India

²Department of Mathematics, University of New Mexico, Gallup, USA

†*Corresponding author*

Abstract

Chakrabarty (1979), Khoshnevisan et al. (2007), Sahai and Ray (1980), Ismail et al. (2011) and Solanki et al. (2012) proposed estimators for estimating population mean \bar{Y} . Up to the first order of approximation and under optimum conditions, the minimum mean squared error (MSE) of all the above estimators is equal to the MSE of the regression estimator. In this paper, we have tried to find out the second order biases and mean square errors of these estimators using information on auxiliary variable based on simple random sampling. Finally, we have compared the performance of these estimators with some numerical illustration.

Keywords: Simple Random Sampling, population mean, study variable, auxiliary variable, exponential ratio type estimator, exponential product estimator, Bias and MSE.

1. Introduction

Let $U = (U_1, U_2, U_3, \dots, U_i, \dots, U_N)$ denotes a finite population of distinct and identifiable units. For estimating the population mean \bar{Y} of a study variable Y , let us consider X be the auxiliary variable that are correlated with study variable Y , taking the corresponding values of the units. Let a sample of size n be drawn from this population using simple random sampling without replacement (SRSWOR) and y_i, x_i ($i=1,2,\dots,n$) are the values of the study variable and auxiliary variable respectively for the i -th unit of the sample.

In sampling theory the use of suitable auxiliary information results in considerable reduction in MSE of the ratio estimators. Many authors suggested estimators using some known population parameters of an auxiliary variable. Upadhyaya and Singh (1999), Singh and Tailor (2003), Kadilar and Cingi (2006), Khoshnevisan et al. (2007), Singh et al. (2007), Singh et al. (2008) and Singh and Kumar (2011) suggested estimators in simple random sampling. Most of the authors discussed the properties of estimators along with their first order bias and MSE. Hossain et al. (2006) studied some estimators in second order approximation. In this study we have studied properties of some estimators under second order of approximation.

2. Some Estimators in Simple Random Sampling

For estimating the population mean \bar{Y} of Y , Chakrabarty (1979) proposed ratio type estimator -

$$t_1 = (1 - \alpha)\bar{y} + \alpha\bar{y}\frac{\bar{X}}{\bar{x}} \quad (2.1)$$

where $\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$ and $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$.

Khoshnevisan et al. (2007) ratio type estimator is given by

$$t_2 = \bar{y} \left[\frac{\bar{X}}{\beta\bar{x} + (1 - \beta)\bar{X}} \right]^g \quad (2.2)$$

where β and g are constants.

Sahai and Ray (1980) proposed an estimator t_3 as

$$t_3 = \bar{y} \left[2 - \left\{ \frac{\bar{x}}{\bar{X}} \right\}^w \right] \quad (2.3)$$

Ismail et al. (2011) proposed an estimator t_4 for estimating the population mean \bar{Y} of Y as

$$t_4 = \bar{y} \left[\frac{\bar{x} + a(\bar{X} - \bar{x})}{\bar{x} + b(\bar{X} - \bar{x})} \right]^p \quad (2.4)$$

where p , a and b are constant.

Also, for estimating the population mean \bar{Y} of Y , Solanki et al. (2012) proposed an estimator t_5 as

$$t_5 = \bar{y} \left[2 - \left\{ \left(\frac{\bar{x}}{\bar{X}} \right)^\lambda \exp \frac{\delta(\bar{x} - \bar{X})}{(\bar{x} + \bar{X})} \right\} \right] \quad (2.5)$$

where λ and δ are constants, suitably chosen by minimizing mean square error of the estimator t_5 .

3. Notations used

Let us define, $e_0 = \frac{\bar{y} - \bar{Y}}{\bar{Y}}$ and $e_1 = \frac{\bar{x} - \bar{X}}{\bar{X}}$, then $E(e_0) = E(e_1) = 0$.

For obtaining the bias and MSE the following lemmas will be used:

Lemma 3.1

- (i) $V(e_0) = E\{(e_0)^2\} = \frac{N-n}{N-1} \frac{1}{n} C_{02} = L_1 C_{02}$
- (ii) $V(e_1) = E\{(e_1)^2\} = \frac{N-n}{N-1} \frac{1}{n} C_{20} = L_1 C_{20}$
- (iii) $COV(e_0, e_1) = E\{(e_0 e_1)\} = \frac{N-n}{N-1} \frac{1}{n} C_{11} = L_1 C_{11}$

Lemma 3.2

- (iv) $E\{(e_1^2 e_0)\} = \frac{(N-n)}{(N-1)} \frac{(N-2n)}{(N-2)} \frac{1}{n^2} C_{21} = L_2 C_{21}$
- (v) $E\{(e_1^3)\} = \frac{(N-n)}{(N-1)} \frac{(N-2n)}{(N-2)} \frac{1}{n^2} C_{30} = L_2 C_{30}$

Lemma 3.3

- (vi) $E(e_1^3 e_0) = L_3 C_{31} + 3L_4 C_{20} C_{11}$
- (vii) $E\{(e_1^4)\} = \frac{(N-n)(N^2 + N - 6nN + 6n^2)}{(N-1)(N-2)(N-3)} \frac{1}{n^3} C_{30} = L_3 C_{40} + 3L_4 C_{20}^2$
- (viii) $E(e_1^2 e_0^2) = L_3 C_{40} + 3L_4 C_{20}$

Where $L_3 = \frac{(N-n)(N^2 + N - 6nN + 6n^2)}{(N-1)(N-2)(N-3)} \frac{1}{n^3}$, $L_4 = \frac{N(N-n)(N-n-1)(n-1)}{(N-1)(N-2)(N-3)} \frac{1}{n^3}$

$$\text{and } C_{pq} = \frac{(X_i - \bar{X})^p (Y_i - \bar{Y})^q}{\bar{X}^p \bar{Y}^q}.$$

Proof of these lemma's are straight forward by using SRSWOR (see Sukhatme and Sukhatme (1970)).

4. First Order Biases and Mean Squared Errors

The expression for the biases of the estimators t_1, t_2, t_3, t_4 and t_5 are respectively given by

$$\text{Bias}(t_1) = \bar{Y} \left[\frac{1}{2} \alpha L_1 C_{20} - \alpha L_1 C_{11} \right] \quad (4.1)$$

$$\text{Bias}(t_2) = \bar{Y} \left[\frac{g(g+1)}{2} L_1 C_{20} - g\beta L_1 C_{11} \right] \quad (4.2)$$

$$\text{Bias}(t_3) = \bar{Y} \left[-\frac{w(w-1)}{2} L_1 C_{20} - w L_1 C_{11} \right] \quad (4.3)$$

$$\text{Bias}(t_4) = \bar{Y} \left[b D L_1 C_{20} - D L_1 C_{11} + \frac{D(b-a)(p-1)}{2} L_1 C_{20} \right] \quad (4.4)$$

$$\text{Bias}(t_5) = \bar{Y} \left[-\frac{K(K-1)}{2} L_1 C_{20} - K L_1 C_{11} \right] \quad (4.5)$$

where, $D = p(b-a)$ and $k = \frac{(\delta + 2\lambda)}{2}$.

Expression for the MSE's of the estimators t_1, t_2, t_3, t_4 and t_5 are, respectively given by

$$\text{MSE}(t_1) = \bar{Y}^2 \left[L_1 C_{02} + \alpha^2 L_1 C_{20} - 2\alpha L_1 C_{11} \right] \quad (4.6)$$

$$\text{MSE}(t_2) = \bar{Y}^2 \left[L_1 C_{02} + g^2 \beta^2 L_1 C_{20} - 2g\beta L_1 C_{11} \right] \quad (4.7)$$

$$\text{MSE}(t_3) = \bar{Y}^2 \left[L_1 C_{02} + w^2 L_1 C_{20} - 2w L_1 C_{11} \right] \quad (4.8)$$

$$\text{MSE}(t_4) = \bar{Y}^2 \left[L_1 C_{02} + D L_1 C_{20} - 2D L_1 C_{11} \right] \quad (4.9)$$

$$\text{MSE}(t_5) = \bar{Y}^2 \left[L_1 C_{02} + k^2 L_1 C_{20} - 2k L_1 C_{11} \right] \quad (4.10)$$

5. Second Order Biases and Mean Squared Errors

Expressing estimator t_i 's ($i=1,2,3,4$) in terms of e 's ($i=0,1$), we get

$$t_1 = \bar{Y}(1 + e_0) \left\{ (1 - \alpha) + \alpha(1 + e_1)^{-1} \right\}$$

Or

$$t_1 - \bar{Y} = \bar{Y} \left\{ e_0 + \frac{e_1}{2} + \frac{\alpha}{2} e_1^2 - \alpha e_0 e_1 + \alpha e_0 e_1^2 - \frac{\alpha}{6} e^3 - \frac{\alpha}{6} e_0 e_1^3 + \frac{\alpha}{24} e^4 \right\} \quad (5.1)$$

Taking expectations, we get the bias of the estimator t_1 up to the second order of approximation as

$$\begin{aligned} \text{Bias}_2(t_1) = \bar{Y} \left[\frac{\alpha}{2} L_1 C_{20} - \alpha L_1 C_{11} - \frac{\alpha}{6} L_2 C_{30} + \alpha L_2 C_{21} - \frac{\alpha}{6} (L_3 C_{31} + 3L_4 C_{20} C_{11}) \right. \\ \left. + \frac{\alpha}{24} (L_3 C_{40} + 3L_4 C_{20}^2) \right] \end{aligned} \quad (5.2)$$

Similarly, we get the biases of the estimator's t_2 , t_3 , t_4 and t_5 up to second order of approximation, respectively as

$$\begin{aligned} \text{Bias}_2(t_2) = \bar{Y} \left[\frac{g(g+1)}{2} \beta^2 L_1 C_{20} - g\beta L_1 C_{11} - \frac{g(g+1)}{2} \beta^2 L_2 C_{21} - \frac{g(g+1)(g+2)}{6} \beta^3 L_2 C_{30} \right. \\ \left. - \frac{g(g+1)(g+2)}{6} \beta^3 (L_3 C_{31} + 3L_4 C_{20} C_{11}) \right. \\ \left. + \frac{g(g+1)(g+2)(g+3)}{24} \beta^4 (L_3 C_{40} + 3L_4 C_{20}^2) \right] \end{aligned} \quad (5.3)$$

$$\begin{aligned} \text{Bias}_2(t_3) = \bar{Y} \left[-\frac{w(w-1)}{2} L_1 C_{20} - w L_1 C_{11} - \frac{w(w-1)}{2} L_2 C_{21} - \frac{w(w-1)(w-2)}{6} L_2 C_{30} \right. \\ \left. - \frac{w(w-1)(w-2)}{6} (L_3 C_{31} + 3L_4 C_{20} C_{11}) \right. \\ \left. - \frac{w(w-1)(w-2)(w-3)}{24} (L_3 C_{40} + 3L_4 C_{20}^2) \right] \end{aligned} \quad (5.4)$$

$$\begin{aligned} \text{Bias}_2(t_4) = \bar{Y} \left[\frac{(D_1 b D)}{2} L_1 C_{20} - D L_1 C_{11} + \frac{(b D + D_1)}{2} L_2 C_{21} - \frac{(b^2 D + 2b D_1 + D_2)}{2} L_2 C_{30} \right. \\ \left. - (b^2 D + 2b D_1) (L_3 C_{31} + 3L_4 C_{20} C_{11}) \right. \\ \left. + \frac{(b^3 D + 3b^2 D_1 + 3b D_2 + D_3)}{2} (L_3 C_{40} + 3L_4 C_{20}^2) \right] \end{aligned} \quad (5.5)$$

$$\text{where, } D_1 = D \frac{(b-a)(p-1)}{2} \quad D_2 = D_1 \frac{(b-a)(p-2)}{3}.$$

$$\begin{aligned} \text{Bias}_2(t_5) = \bar{Y} \left[-\frac{k(k-1)}{2} L_1 C_{20} - k L_1 C_{11} - \frac{k(k-1)}{2} L_2 C_{21} - M L_2 C_{30} - M(L_3 C_{31} + 3L_4 C_{20} C_{11}) \right. \\ \left. - N(L_3 C_{40} + 3L_4 C_{20}^2) \right] \end{aligned} \quad (5.6)$$

$$\text{Where, } M = \frac{1}{2} \left\{ \frac{(\delta^3 - 6\delta^2)}{24} + \frac{(\alpha(\delta^2 - 2\delta))}{4} + \frac{\lambda(\lambda-1)}{2} \delta + \frac{\lambda(\lambda-1)(\lambda-2)}{3} \right\},$$

$$k = \frac{(\delta + 2\lambda)}{2},$$

$$N = \frac{1}{8} \left\{ \frac{(\delta^4 - 12\delta^3 + 12\delta^2)}{48} + \frac{(\alpha(\delta^3 - 6\delta))}{6} + \frac{\lambda(\lambda-1)}{2} (\delta^2 - 2\delta) + \frac{\lambda(\lambda-1)(\lambda-2)(\lambda-3)}{3} \right\}.$$

The MSE's of the estimators t_1, t_2, t_3, t_4 and t_5 up to the second order of approximation are, respectively given by

$$\begin{aligned} \text{MSE}_2(t_1) = \bar{Y}^2 \left[L_1 C_{02} + \alpha^2 L_1 C_{20} - 2\alpha L_1 C_{11} - \alpha^2 L_2 C_{30} + (2\alpha^2 + \alpha) L_2 C_{21} \right. \\ \left. - 2\alpha^2 (L_3 C_{31} + 3L_4 C_{20} C_{11}) \right. \\ \left. + \alpha(\alpha+1) (L_3 C_{22} + 3L_4 (C_{20} C_{02} + C_{11}^2)) + \frac{5}{24} \alpha^2 (L_3 C_{40} + 3L_4 C_{20}^2) \right] \end{aligned} \quad (5.7)$$

$$\begin{aligned} \text{MSE}_2(t_2) = \bar{Y}^2 \left[L_1 C_{02} + g^2 \beta^2 L_1 C_{20} - 2\beta g L_1 C_{11} - \beta^3 g^2 (g+1) L_2 C_{30} + g(3g+1) \beta^2 L_2 C_{21} \right. \\ \left. - 2\beta g L_2 C_{12} - \left\{ \frac{7g^3 + 9g^2 + 2g}{3} \right\} \beta^3 (L_3 C_{31} + 3L_4 C_{20} C_{11}) \right. \\ \left. + g(2g+1) \beta^2 (L_3 C_{22} + 3L_4 (C_{20} C_{02} + C_{11}^2)) \right. \\ \left. + \left\{ \frac{2g^3 + 9g^2 + 10g + 3}{6} \right\} \beta^4 (L_3 C_{40} + 3L_4 C_{20}^2) \right] \end{aligned} \quad (5.8)$$

$$\begin{aligned}
\text{MSE}_2(t_3) = & \bar{Y}^2 \left[L_1 C_{02} + w^2 L_1 C_{20} - 2w L_1 C_{11} - w^2 (w-1) L_2 C_{30} + w(w+1) L_2 C_{21} - 2w L_2 C_{12} \right. \\
& + \left. \left\{ \frac{5w^3 - 3w^2 - 2w}{3} \right\} (L_3 C_{31} + 3L_4 C_{20} C_{11}) \right. \\
& + \left. w(L_3 C_{22} + 3L_4 (C_{20} C_{02} + C_{11}^2)) + \left\{ \frac{7w^4 - 18w^3 + 11w^2}{24} \right\} (L_3 C_{40} + 3L_4 C_{20}^2) \right]
\end{aligned} \tag{5.9}$$

$$\begin{aligned}
\text{MSE}_2(t_4) = & \bar{Y}^2 \left[L_1 C_{02} + D^2 L_1 C_{20} - 2D L_1 C_{11} - 4DD_1 L_2 C_{30} + (2bD + 2D_1 + 2D^2) L_2 C_{21} - 2D L_2 C_{12} \right. \\
& + \left. \{ 2D^2 + 2b^2 D + 2DD_1 + 4bD_1 + 4bD^2 \} (L_3 C_{31} + 3L_4 C_{20} C_{11}) \right. \\
& + \left. \{ D^2 + 2D_1 + 2bD \} (L_3 C_{22} + 3L_4 (C_{20} C_{02} + C_{11}^2)) \right. \\
& + \left. \{ 3b^2 D^2 + D_1^2 + 2DD_2 + 12bDD_1 \} (L_3 C_{40} + 3L_4 C_{20}^2) \right]
\end{aligned} \tag{5.10}$$

$$\begin{aligned}
\text{MSE}_2(t_5) = & \bar{Y}^2 \left[L_1 C_{02} + k^2 L_1 C_{20} - 2k L_1 C_{11} + k L_2 C_{21} - 2k L_2 C_{12} + k^2 (k-1) L_2 C_{30} \right. \\
& + 2k^2 (k-1) (L_3 C_{31} + 3L_4 C_{20} C_{11}) + k (L_3 C_{22} + 3L_4 (C_{20} C_{02} + C_{11}^2)) \\
& + \left. \frac{(k^2 - k)^2}{4} (L_3 C_{40} + 3L_4 C_{20}^2) \right]
\end{aligned} \tag{5.11}$$

6. Numerical Illustration

For a natural population data, we have calculated the biases and the mean square error's of the estimator's and compare these biases and MSE's of the estimator's under first and second order of approximations.

Data Set

The data for the empirical analysis are taken from 1981, Utter Pradesh District Census Handbook, Aligar. The population consist of 340 villages under koil police station, with Y=Number of agricultural labour in 1981 and X=Area of the villages (in acre) in 1981. The following values are obtained

$\bar{Y} = 73.76765$, $\bar{X} = 2419.04$, $N = 340$, $n = 70$, $n' = 120$, $n=70$, $C_{02}=0.7614$, $C_{11}=0.2667$,
 $C_{03}=2.6942$, $C_{12}=0.0747$, $C_{12}=0.1589$, $C_{30}=0.7877$, $C_{13}=0.1321$, $C_{31}=0.8851$, $C_{04}=17.4275$
 $C_{22}=0.8424$, $C_{40}=1.3051$

Table 6.1: Biases and MSE's of the estimators

Estimator	Bias		MSE	
	First order	Second order	First order	Second order
t_1	0.0044915	0.004424	39.217225	39.45222
t_2	0	-0.00036	39.217225 (for $g=1$)	39.33552 (for $g=1$)
t_3	-0.04922	-0.04935	39.217225	39.29102
t_4	0.2809243	-0.60428	39.217225	39.44855
t_5	-0.027679	-0.04911	39.217225	39.27187

In the Table 6.1 the biases and MSE's of the estimators t_1 , t_2 , t_3 , t_4 and t_5 are written under first order and second order of approximations. For all the estimators t_1 , t_2 , t_3 , t_4 and t_5 , it was observed that the value of the biases decreased and the value of the MSE's increased for second order approximation. MSE's of the estimators up to the first order of approximation under optimum conditions are same. From Table 6.1 we observe that under

second order of approximation the estimator t_5 is best followed by t_3 , and t_2 among the estimators considered here for the given data set.

7. Estimators under stratified random sampling

In survey sampling, it is well established that the use of auxiliary information results in substantial gain in efficiency over the estimators which do not use such information. However, in planning surveys, the stratified sampling has often proved needful in improving the precision of estimates over simple random sampling. Assume that the population U consist of L strata as $U=U_1, U_2, \dots, U_L$. Here the size of the stratum U_h is N_h , and the size of simple random sample in stratum U_h is n_h , where $h=1, 2, \dots, L$.

The Chakrabarty(1979) ratio- type estimator under stratified random sampling is given by

$$t'_1 = (1 - \alpha)\bar{y}_{st} + \alpha\bar{y}_{st} \frac{\bar{X}}{\bar{X}_{st}} \quad (7.1)$$

where,

$$\bar{y}_h = \frac{1}{n_h} \sum_{i=1}^{n_h} y_{hi}, \quad \bar{x}_h = \frac{1}{n_h} \sum_{i=1}^{n_h} x_{hi},$$

$$\bar{y}_{st} = \sum_{h=1}^L w_h \bar{y}_h, \quad \bar{x}_{st} = \sum_{h=1}^L w_h \bar{x}_h, \quad \text{and} \quad \bar{X} = \sum_{h=1}^L w_h \bar{X}_h.$$

Khoshnevisan et al. (2007) ratio- type estimator under stratified random sampling is given by

$$t'_2 = \bar{y}_{st} \left[\frac{\bar{X}}{\beta \bar{x}_{st} + (1 - \beta)\bar{X}} \right]^g \quad (7.2)$$

where g is a constant, for $g=1$, t'_2 is same as conventional ratio estimator whereas for $g = -1$, it becomes conventional product type estimator.

Sahai and Ray (1980) estimator t_3 under stratified random sampling is given by

$$t'_3 = \bar{y}_{st} \left[2 - \left\{ \frac{\bar{x}_{st}}{\bar{X}} \right\}^w \right] \quad (7.3)$$

Ismail et al. (2011) estimator under stratified random sampling t'_4 is given by

$$t'_4 = \bar{y}_{st} \left[\frac{\bar{x} + a(\bar{X} - \bar{x}_{st})}{\bar{x} + b(\bar{X} - \bar{x}_{st})} \right]^p \quad (7.4)$$

Solanki et al. (2012) estimator under stratified random sampling is given as

$$t'_5 = \bar{y}_{st} \left[2 - \left\{ \left(\frac{\bar{x}_{st}}{\bar{X}} \right)^\lambda \exp \frac{\delta(\bar{x}_{st} - \bar{X})}{(\bar{x}_{st} + \bar{X})} \right\} \right] \quad (7.5)$$

where λ and δ are the constants, suitably chosen by minimizing MSE of the estimator t'_5 .

8. Notations used under stratified random sampling

Let us define, $e_0 = \frac{\bar{y}_{st} - \bar{Y}}{\bar{Y}}$ and $e_1 = \frac{\bar{x}_{st} - \bar{X}}{\bar{X}}$, then $E(e_0) = E(e_1) = 0$.

To obtain the bias and MSE of the proposed estimators, we use the following notations in the rest of the article:

$$\bar{y}_{st} = \sum_{h=1}^L w_h \bar{y}_h = \bar{Y}(1 + e_0),$$

$$\bar{x}_{st} = \sum_{h=1}^L w_h \bar{x}_h = \bar{X}(1 + e_1),$$

such that,

$$E(e_0) = E(e_1) = E(e_2) = 0, \text{ and}$$

$$V_{rs} = \sum_{h=1}^L W_h^{r+s} E \left[(\bar{x}_h - \bar{X}_h)^r (\bar{y}_h - \bar{Y}_h)^s \right]$$

Also

$$E(e_0^2) = \frac{\sum_{h=1}^L w_h^2 \gamma_h S_{yh}^2}{\bar{Y}^2} = V_{20}$$

$$E(e_1^2) = \frac{\sum_{h=1}^L w_h^2 \gamma_h S_{xyh}^2}{\bar{X}^2} = V_{02}$$

$$E(e_0 e_1) = \frac{\sum_{h=1}^L w_h^2 \gamma_h S_{xyh}^2}{\bar{Y}\bar{X}} = V_{11}$$

Where

$$S_{yh}^2 = \frac{\sum_{i=1}^{N_h} (\bar{y}_h - \bar{Y}_h)^2}{N_h - 1}, \quad S_{xh}^2 = \frac{\sum_{i=1}^{N_h} (\bar{x}_h - \bar{X}_h)^2}{N_h - 1}, \quad S_{xyh} = \frac{\sum_{i=1}^{N_h} (\bar{x}_h - \bar{X}_h)(\bar{y}_h - \bar{Y}_h)}{N_h - 1}$$

$$\gamma_h = \frac{1 - f_h}{n_h}, \quad f_h = \frac{n_h}{N_h}, \quad w_h = \frac{N_h}{n_h}.$$

Some additional notations for second order approximation,

$$V_{rs} = \sum_{h=1}^L W_h^{r+s} \frac{1}{\bar{Y}^r \bar{X}^s} E[(\bar{y}_h - \bar{Y}_h)^s (\bar{x}_h - \bar{X}_h)^r]$$

Where,

$$C_{rs(h)} = \frac{1}{N_h} \sum_{i=1}^{N_h} [(\bar{y}_h - \bar{Y}_h)^s (\bar{x}_h - \bar{X}_h)^r]$$

$$V_{12} = \sum_{h=1}^L W_h^3 \frac{k_{1(h)} C_{12(h)}}{\bar{Y}\bar{X}^2} \quad V_{21} = \sum_{h=1}^L W_h^3 \frac{k_{1(h)} C_{21(h)}}{\bar{Y}^2 \bar{X}} \quad V_{30} = \sum_{h=1}^L W_h^3 \frac{k_{1(h)} C_{30(h)}}{\bar{Y}^3}$$

$$V_{03} = \sum_{h=1}^L W_h^3 \frac{k_{1(h)} C_{03(h)}}{\bar{X}^3} \quad V_{13} = \sum_{h=1}^L W_h^4 \frac{k_{2(h)} C_{13(h)} + 3k_{3(h)} C_{01(h)} C_{02(h)}}{\bar{Y}\bar{X}^3}$$

$$V_{04} = \sum_{h=1}^L W_h^4 \frac{k_{2(h)} C_{04(h)} + 3k_{3(h)} C_{02(h)}^2}{\bar{X}^4}$$

$$V_{22} = \sum_{h=1}^L W_h^4 \frac{k_{2(h)} C_{22(h)} + k_{3(h)} (C_{01(h)} C_{02(h)} + 2C_{11(h)}^2)}{\bar{Y}^2 \bar{X}^2}$$

Where

$$k_{1(h)} = \frac{(N_h - n_h)(N_h - 2n_h)}{n^2(N_h - 1)(N_h - 2)}$$

$$k_{2(h)} = \frac{(N_h - n_h)(N_h + 1)N_h - 6n_h(N_h - n_h)}{n^3(N_h - 1)(N_h - 2)(N_h - 3)}$$

$$k_{3(h)} = \frac{(N_h - n_h)N_h(N_h - n_h - 1)(n_h - 1)}{n^3(N_h - 1)(N_h - 2)(N_h - 3)}$$

9. First Order Biases and Mean Squared Errors

The biases of the estimators t'_1, t'_2, t'_3, t'_4 and t'_5 are respectively given by

$$\text{Bias}(t'_1) = \bar{Y} \left[\frac{1}{2} \alpha L_1 V_{02} - \alpha V_{11} \right] \quad (9.1)$$

$$\text{Bias}(t'_2) = \bar{Y} \left[\frac{g(g+1)}{2} \beta^2 V_{02} - g\beta V_{11} \right] \quad (9.2)$$

$$\text{Bias}(t'_3) = \bar{Y} \left[-\frac{w(w-1)}{2} V_{02} - wV_{11} \right] \quad (9.3)$$

$$\text{Bias}(t'_4) = \bar{Y} \left[bDV_{02} - DV_{11} + \frac{D(b-a)(p-1)}{2} V_{02} \right] \quad (9.4)$$

$$\text{Bias}(t'_5) = \bar{Y} \left[-\frac{K(K-1)}{2} V_{02} - KV_{11} \right] \quad (9.5)$$

Where, $D = p(b-a)$ and $k = \frac{(\delta + 2\lambda)}{2}$.

The MSE's of the estimators t'_1, t'_2, t'_3, t'_4 and t'_5 are respectively given by

$$\text{MSE}(t'_1) = \bar{Y}^2 [V_{20} + \alpha^2 V_{02} - 2\alpha V_{11}] \quad (9.6)$$

$$\text{MSE}(t'_2) = \bar{Y}^2 [V_{20} + g^2 \beta^2 V_{02} - 2g\beta V_{11}] \quad (9.7)$$

$$\text{MSE}(t'_3) = \bar{Y}^2 [V_{20} + w^2 V_{02} - 2wV_{11}] \quad (9.8)$$

$$\text{MSE}(t'_4) = \bar{Y}^2 [V_{20} + DV_{02} - 2DV_{11}] \quad (9.9)$$

$$\text{MSE}(t'_5) = \bar{Y}^2 [V_{20} + k^2 V_{02} - 2kV_{11}] \quad (9.10)$$

10. Second Order Biases and Mean Squared Errors

Expressing estimator t_i 's ($i=1,2,3,4$) in terms of e 's ($i=0,1$), we get

$$t'_1 = \bar{Y}(1 + e_0) \left\{ (1 - \alpha) + \alpha(1 + e_1)^{-1} \right\}$$

Or

$$t'_1 - \bar{Y} = \bar{Y} \left\{ e_0 + \frac{e_1}{2} + \frac{\alpha}{2} e_1^2 - \alpha e_0 e_1 + \alpha e_0 e_1^2 - \frac{\alpha}{6} e^3 - \frac{\alpha}{6} e_0 e_1^3 + \frac{\alpha}{24} e^4 \right\} \quad (10.1)$$

Taking expectations, we get the bias of the estimator t'_1 up to the second order of approximation as

$$\text{Bias}_2(t'_1) = \bar{Y} \left[\frac{\alpha}{2} V_{02} - \alpha V_{11} - \frac{\alpha}{6} V_{03} + \alpha V_{12} - \frac{\alpha}{6} V_{13} + \frac{\alpha}{24} V_{04} \right] \quad (10.2)$$

Similarly we get the Biases of the estimator's t'_2, t'_3, t'_4 and t'_5 up to second order of approximation, respectively as

$$\text{Bias}_2(t'_2) = \bar{Y} \left[\frac{g(g+1)}{2} \beta^2 V_{02} - g\beta V_{11} - \frac{g(g+1)}{2} \beta^2 V_{12} - \frac{g(g+1)(g+2)}{6} \beta^3 V_{30} \right. \\ \left. - \frac{g(g+1)(g+2)}{6} \beta^3 V_{31} + \frac{g(g+1)(g+2)(g+3)}{24} \beta^4 V_{04} \right] \quad (10.3)$$

$$\text{Bias}_2(t'_3) = \bar{Y} \left[-\frac{w(w-1)}{2} V_{02} - wV_{11} - \frac{w(w-1)}{2} V_{12} - \frac{w(w-1)(w-2)}{6} V_{03} \right. \\ \left. - \frac{w(w-1)(w-2)}{6} V_{31} - \frac{w(w-1)(w-2)(w-3)}{24} V_{04} \right] \quad (10.4)$$

$$\text{Bias}_2(t'_4) = \bar{Y} \left[\frac{(D_1 b D)}{2} V_{02} - D V_{11} + (bD + D_1) V_{12} - (b^2 D + 2bD_1 + D_2) V_{03} \right. \\ \left. - (b^2 D + 2bD_1) V_{31} + (b^3 D + 3b^2 D_1 + 3bD_2 + D_3) V_{04} \right] \quad (10.5)$$

$$\text{Where, } D = p(b-a) \quad D_1 = D \frac{(b-a)(p-1)}{2} \quad D_2 = D_1 \frac{(b-a)(p-2)}{3}$$

$$\text{Bias}_2(t'_5) = \bar{Y} \left[-\frac{k(k-1)}{2} V_{02} - kV_{11} - \frac{k(k-1)}{2} V_{12} - M V_{03} - M V_{31} - N V_{04} \right] \quad (10.6)$$

$$\text{Where, } M = \frac{1}{2} \left\{ \frac{(\delta^3 - 6\delta^2)}{24} + \frac{(\alpha(\delta^2 - 2\delta))}{4} + \frac{\lambda(\lambda - 1)}{2} \delta + \frac{\lambda(\lambda - 1)(\lambda - 2)}{3} \right\}, \quad k = \frac{(\delta + 2\lambda)}{2}$$

$$N = \frac{1}{8} \left\{ \frac{(\delta^4 - 12\delta^3 + 12\delta^2)}{48} + \frac{(\alpha(\delta^3 - 6\delta))}{6} + \frac{\lambda(\lambda - 1)}{2} (\delta^2 - 2\delta) + \frac{\lambda(\lambda - 1)(\lambda - 2)(\lambda - 3)}{3} \right\}$$

Following are the MSE of the estimators t'_1, t'_2, t'_3, t'_4 and t'_5 up to second order of approximation

$$\begin{aligned} \text{MSE}_2(t'_1) = \bar{Y}^2 & \left[V_{20} + \alpha^2 V_{02} - 2\alpha V_{11} - \alpha^2 V_{03} + (2\alpha^2 + \alpha) V_{12} \right. \\ & \left. - 2\alpha^2 V_{31} + \alpha(\alpha + 1) V_{22} + \frac{5}{24} \alpha^2 V_{04} \right] \end{aligned} \quad (10.7)$$

$$\begin{aligned} \text{MSE}_2(t'_2) = \bar{Y}^2 & \left[V_{20} + g^2 \beta^2 V_{02} - 2\beta g V_{11} - \beta^3 g^2 (g + 1) V_{03} + g(3g + 1) \beta^2 V_{12} \right. \\ & \left. - 2\beta g V_{21} - \left\{ \frac{7g^3 + 9g^2 + 2g}{3} \right\} \beta^3 V_{31} + g(2g + 1) \beta^2 V_{22} \right. \\ & \left. + \left\{ \frac{2g^3 + 9g^2 + 10g + 3}{6} \right\} \beta^4 V_{04} \right] \end{aligned} \quad (10.8)$$

$$\begin{aligned} \text{MSE}_2(t'_3) = \bar{Y}^2 & \left[V_{20} + w^2 V_{02} - 2w V_{11} - w^2 (w - 1) V_{03} + w(w + 1) V_{12} - 2w V_{21} \right. \\ & \left. + \left\{ \frac{5w^3 - 3w^2 - 2w}{3} \right\} V_{31} + w V_{22} + \left\{ \frac{7w^4 - 18w^3 + 11w^2}{24} \right\} V_{04} \right] \end{aligned} \quad (10.9)$$

$$\begin{aligned} \text{MSE}_2(t'_4) = \bar{Y}^2 & \left[V_{20} + D^2 V_{02} - 2D V_{11} - 4DD_1 V_{03} + (2bD + 2D_1 + 2D^2) V_{12} - 2D V_{21} \right. \\ & \left. + \left\{ 2D^2 + 2b^2 D + 2DD_1 + 4bD_1 + 4bD^2 \right\} V_{31} + \left\{ D^2 + 2D_1 + 2bD \right\} V_{22} \right. \\ & \left. + \left\{ 3b^2 D^2 + D_1^2 + 2DD_2 + 12bDD_1 \right\} V_{04} \right] \end{aligned} \quad (10.10)$$

$$\begin{aligned} \text{MSE}_2(t'_5) = \bar{Y}^2 & \left[V_{20} + k^2 V_{02} - 2k V_{11} + k V_{12} - 2k V_{21} + k^2 (k - 1) V_{03} \right. \\ & \left. + 2k^2 (k - 1) V_{31} + k V_{22} + \frac{(k^2 - k)^2}{4} V_{04} \right] \end{aligned} \quad (10.11)$$

11. Numerical Illustration

For the natural population data, we shall calculate the bias and the mean square error of the estimator and compare Bias and MSE for the first and second order of approximation.

Data Set-1

To illustrate the performance of above estimators, we have considered the natural Data given in *Singh and Chaudhary (1986, p.162)*.

The data were collected in a pilot survey for estimating the extent of cultivation and production of fresh fruits in three districts of Uttar- Pradesh in the year 1976-1977.

Table 11.1: Biases and MSE's of the estimators

Estimator	Bias		MSE	
	First order	Second order	First order	second order
t'_1	-10.82707903	-13.65734654	1299.110219	1372.906438
t'_2	-10.82707903	6.543275811	1299.110219	1367.548263
t'_3	-27.05776113	-27.0653128	1299.110219	1417.2785
t'_4	11.69553975	-41.84516913	1299.110219	2605.736045
t'_5	-22.38574093	-14.95110301	1299.110219	2440.644397

From Table 11.1 we observe that the MSE's of the estimators t'_1, t'_2, t'_3, t'_4 and t'_5 are same up to the first order of approximation but the biases are different. The MSE of the estimator t'_2 is minimum under second order of approximation followed by the estimator t'_1 and other estimators.

Conclusion

In this study we have considered some estimators whose MSE's are same up to the first order of approximation. We have extended the study to second order of approximation to search for best estimator in the sense of minimum variance. The properties of the estimators are studied under SRSWOR and stratified random sampling. We have observed from Table 6.1 and Table 11.1 that the behavior of the estimators changes dramatically when we consider the terms up to second order of approximation.

REFERENCES

- Bahl, S. and Tuteja, R.K. (1991) : Ratio and product type exponential estimator. Information and Optimization Science XIII 159-163.
- Chakrabarty, R.P. (1979) : Some ratio estimators, Journal of the Indian Society of Agricultural Statistics **31**(1), 49–57.
- Hossain, M.I., Rahman, M.I. and Tareq, M. (2006) : Second order biases and mean squared errors of some estimators using auxiliary variable. SSRN.
- Ismail, M., Shahbaz, M.Q. and Hanif, M. (2011) : A general class of estimator of population mean in presence of non–response. Pak. J. Statist. 27(4), 467-476.
- Khoshnevisan, M., Singh, R., Chauhan, P., Sawan, N., and Smarandache, F. (2007). A general family of estimators for estimating population mean using known value of some population parameter(s), Far East Journal of Theoretical Statistics 22 181–191.
- Ray, S.K. and Sahai, A (1980) : Efficient families of ratio and product type estimators, Biometrika **67**(1) , 211–215.
- Singh, D. and Chudhary, F.S. (1986): *Theory and analysis of sample survey designs*. Wiley Eastern Limited, New Delhi.

- Singh, H.P. and Tailor, R. (2003). Use of known correlation coefficient in estimating the finite population mean. *Statistics in Transition* 6, 555-560.
- Singh, R., Chauhan, P., Sawan, N., and Smarandache, F. (2007): *Auxiliary Information and A Priori Values in Construction of Improved Estimators*. Renaissance High Press.
- Singh, R., Chauhan, P. and Sawan, N. (2008): On linear combination of Ratio-product type exponential estimator for estimating finite population mean. *Statistics in Transition*,9(1),105-115.
- Singh, R., Kumar, M. and Smarandache, F. (2008): Almost Unbiased Estimator for Estimating Population Mean Using Known Value of Some Population Parameter(s). *Pak. J. Stat. Oper. Res.*, 4(2) pp63-76.
- Singh, R. and Kumar, M. (2011): A note on transformations on auxiliary variable in survey sampling. *MASA*, 6:1, 17-19.
- Solanki, R.S., Singh, H. P. and Rathour, A. (2012) : An alternative estimator for estimating the finite population mean using auxiliary information in sample surveys. *ISRN Probability and Statistics* doi:10.5402/2012/657682
- Srivastava, S.K. (1967) : An estimator using auxiliary information in sample surveys. *Cal. Stat. Ass. Bull.* 15:127-134.
- Sukhatme, P.V. and Sukhatme, B.V. (1970): *Sampling theory of surveys with applications*. Iowa State University Press, Ames, U.S.A.
- Upadhyaya, L. N. and Singh, H. P. (199): Use of transformed auxiliary variable in estimating the finite population mean. *Biom. Jour.*, 41, 627-636.

IMPROVEMENT IN ESTIMATING THE POPULATION MEAN USING DUAL TO RATIO-CUM-PRODUCT ESTIMATOR IN SIMPLE RANDOM SAMPLING

¹Olufadi Yunusa, ²†Rajesh Singh and ³Florentin Smarandache

¹Department of Statistics and Mathematical Sciences
Kwara State University, P.M.B 1530, Malete, Nigeria

²Department of Statistics, Banaras Hindu University, Varanasi (U.P.), India

³Department of Mathematics, University of New Mexico, Gallup, USA

†Corresponding author

ABSTRACT

In this paper, we propose a new estimator for estimating the finite population mean using two auxiliary variables. The expressions for the bias and mean square error of the suggested estimator have been obtained to the first degree of approximation and some estimators are shown to be a particular member of this estimator. Furthermore, comparison of the suggested estimator with the usual unbiased estimator and other estimators considered in this paper is carried out. In addition, an empirical study with two natural data from literature is used to expound the performance of the proposed estimator with respect to others.

Keywords: Dual-to-ratio estimator; finite population mean; mean square error; multi-auxiliary variable; percent relative efficiency; ratio-cum-product estimator

1. INTRODUCTION

It is well known that the use of auxiliary information in sample survey design results in efficient estimate of population parameters (e.g. mean) under some realistic conditions. This information may be used at the design stage (leading, for instance, to stratification,

systematic or probability proportional to size sampling designs), at the estimation stage or at both stages. The literature on survey sampling describes a great variety of techniques for using auxiliary information by means of ratio, product and regression methods. Ratio and product type estimators take advantage of the correlation between the auxiliary variable, x and the study variable, y . For example, when information is available on the auxiliary variable that is positively (high) correlated with the study variable, the ratio method of estimation is a suitable estimator to estimate the population mean and when the correlation is negative the product method of estimation as envisaged by Robson (1957) and Murthy (1964) is appropriate.

Quite often information on many auxiliary variables is available in the survey which can be utilized to increase the precision of the estimate. In this situation, Olkin (1958) was the first author to deal with the problem of estimating the mean of a survey variable when auxiliary variables are made available. He suggested the use of information on more than one supplementary characteristic, positively correlated with the study variable, considering a linear combination of ratio estimators based on each auxiliary variable separately. The coefficients of the linear combination were determined so as to minimize the variance of the estimator. Analogously to Olkin, Singh (1967) gave a multivariate expression of Murthy's (1964) product estimator, while Raj (1965) suggested a method for using multi-auxiliary variables through a linear combination of single difference estimators. More recently, Abu-Dayyeh et al. (2003), Kadilar and Cingi (2004, 2005), Perri (2004, 2005), Dianna and Perri (2007), Malik and Singh (2012) among others have suggested estimators for \bar{Y} using information on several auxiliary variables.

Motivated by Srivenkataramana (1980), Bandyopadhyay (1980) and Singh et al. (2005) and with the aim of providing a more efficient estimator; we propose, in this paper, a new estimator for \bar{Y} when two auxiliary variables are available.

2. BACKGROUND TO THE SUGGESTED ESTIMATOR

Consider a finite population $P = (P_1, P_2, \dots, P_N)$ of N units. Let a sample s of size n be drawn from this population by simple random sampling without replacements (SRSWOR). Let y_i and (x_i, z_i) represents the value of a response variable y and two auxiliary variables (x, z) are available. The units of this finite population are identifiable in the sense that they are uniquely labeled from 1 to N and the label on each unit is known. Further, suppose in a survey problem, we are interested in estimating the population mean \bar{Y} of y , assuming that the population means (\bar{X}, \bar{Z}) of (x, z) are known. The traditional ratio and product estimators for \bar{Y} are given as

$$\bar{y}_R = \bar{y} \frac{\bar{X}}{\bar{x}} \text{ and } \bar{y}_P = \bar{y} \frac{\bar{z}}{\bar{Z}}$$

respectively, where $\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$, $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$ and $\bar{z} = \frac{1}{n} \sum_{i=1}^n z_i$ are the sample means of y , x and z respectively.

Singh (1969) improved the ratio and product method of estimation given above and suggested the “ratio-cum-product” estimator for \bar{Y} as $\bar{y}_S = \bar{y} \frac{\bar{X}}{\bar{x}} \frac{\bar{z}}{\bar{Z}}$

In literature, it has been shown by various authors; see for example, Reddy (1974) and Srivenkataramana (1978) that the bias and the mean square error of the ratio estimator \bar{y}_R , can be reduced with the application of transformation on the auxiliary variable x . Thus, authors like, Srivenkataramana (1980), Bandyopadhyay (1980) Tracy et al. (1996), Singh et al. (1998), Singh et al. (2005), Singh et al. (2007), Bartkus and Plikusas (2009) and Singh et al. (2011) have improved on the ratio, product and ratio-cum-product method of estimation using the transformation on the auxiliary information. We give below the transformations employed by these authors:

$$x_i^* = (1 + g)\bar{X} - gx_i \text{ and } z_i^* = (1 + g)\bar{Z} - gz_i, \text{ for } i = 1, 2, \dots, N, \quad (1)$$

where $g = \frac{n}{N-n}$.

Then clearly, $\bar{x}^* = (1 + g)\bar{X} - g\bar{x}$ and $\bar{z}^* = (1 + g)\bar{Z} - g\bar{z}$ are also unbiased estimate of \bar{X} and \bar{Z} respectively and $Corr(\bar{y}, \bar{x}^*) = -\rho_{yx}$ and $Corr(\bar{y}, \bar{z}^*) = -\rho_{yz}$. It is to be noted that by using the transformation above, the construction of the estimators for \bar{Y} requires the knowledge of unknown parameters, which restrict the applicability of these estimators. To overcome this restriction, in practice, information on these parameters can be obtained approximately from either past experience or pilot sample survey, inexpensively.

The following estimators \bar{y}_R^* , \bar{y}_P^* and \bar{y}_{SE} are referred to as dual to ratio, product and ratio-cum-product estimators and are due to Srivenkataramana (1980), Bandyopadhyay (1980) and Singh et al. (2005) respectively. They are as presented: $\bar{y}_R^* = \bar{y} \frac{\bar{x}^*}{\bar{X}}$, $\bar{y}_P^* = \bar{y} \frac{\bar{Z}}{\bar{z}^*}$ and $\bar{y}_{SE} = \bar{y} \frac{\bar{x}^*}{\bar{X}} \frac{\bar{Z}}{\bar{z}^*}$

It is well known that the variance of the simple mean estimator \bar{y} , under SRSWOR design is

$$V(\bar{y}) = \lambda S_y^2$$

and to the first order of approximation, the Mean Square Errors (MSE) of \bar{y}_R , \bar{y}_P , \bar{y}_S , \bar{y}_R^* ,

\bar{y}_P^* and \bar{y}_{SE} are, respectively, given by

$$MSE(\bar{y}_R) = \lambda(S_y^2 + R_1^2 S_x^2 - 2R_1 S_{yx})$$

$$MSE(\bar{y}_P) = \lambda(S_y^2 + R_2^2 S_z^2 + 2R_2 S_{yz})$$

$$MSE(\bar{y}_S) = \lambda[S_y^2 - 2D + C]$$

$$MSE(\bar{y}_R^*) = \lambda(S_y^2 + g^2 R_1^2 S_x^2 - 2gR_1 S_{yx})$$

$$MSE(\bar{y}_P^*) = \lambda(S_y^2 + g^2 R_2^2 S_z^2 + 2gR_2 S_{yz})$$

$$MSE(\bar{y}_{SE}) = \lambda(S_y^2 + g^2C - 2gD)$$

where,

$$\lambda = \frac{1-f}{n}, \quad f = \frac{n}{N}, \quad S_y^2 = \frac{1}{N} \sum_{i=1}^N (y_i - \bar{Y})^2, \quad S_{yx} = \frac{1}{N} \sum_{i=1}^N (y_i - \bar{Y})(x_i - \bar{X}), \quad \rho_{yx} = \frac{S_{yx}}{S_y S_x},$$

$$R_1 = \frac{\bar{Y}}{\bar{X}}, \quad R_2 = \frac{\bar{Y}}{\bar{Z}}, \quad C = R_1^2 S_x^2 - 2R_1 R_2 S_{zx} + R_2^2 S_z^2, \quad D = R_1 S_{yx} - R_2 S_{yz} \quad \text{and} \quad S_j^2 \quad \text{for}$$

($j = x, y, z$) represents the variances of x , y and z respectively; while S_{yx} , S_{yz} and S_{zx} denote the covariance between y and x , y and z and z and x respectively. Note that ρ_{yz} , ρ_{zx} , S_x^2 , S_z^2 , S_{yz} and S_{zx} are defined analogously and respective to the subscripts used.

More recently, Sharma and Tailor (2010) proposed a new ratio-cum-dual to ratio estimator of finite population mean in simple random sampling, their estimator with its MSE are respectively given as,

$$\bar{y}_{ST} = \bar{y} \left[\alpha \left(\frac{\bar{X}}{\bar{x}} \right) + (1-\alpha) \left(\frac{\bar{x}^*}{\bar{X}} \right) \right]$$

$$MSE(\bar{y}_{ST}) = \lambda S_y^2 (1 - \rho_{yx}^2).$$

3. PROPOSED DUAL TO RATIO-CUM-PRODUCT ESTIMATOR

Using the transformation given in (1), we suggest a new estimator for \bar{Y} as follows:

$$\bar{y}_{PR} = \bar{y} \left[\theta \left(\frac{\bar{x}^* \bar{Z}}{\bar{X} \bar{z}^*} \right) + (1-\theta) \left(\frac{\bar{X} \bar{z}^*}{\bar{x}^* \bar{Z}} \right) \right]$$

We note that when information on the auxiliary variable z is not used (or variable z takes the value 'unity') and $\theta = 1$, the suggested estimator \bar{y}_{PR} reduces to the 'dual to ratio' estimator \bar{y}_R^* proposed by Srivenkataramana (1980). Also, \bar{y}_{PR} reduces to the 'dual to product' estimator \bar{y}_P^* proposed by Bandyopadhyay (1980) estimator if the information on the auxiliary variate x is not used and $\theta = 0$. Furthermore, the suggested estimator reduces

to the dual to ratio-cum-product estimator suggested by Singh et al. (2005) when $\theta = 1$ and information on the two auxiliary variables x and z are been utilized.

In order to study the properties of the suggested estimator \bar{y}_{PR} (e.g. MSE), we write

$$\bar{y} = \bar{Y}(1 + k_0); \quad \bar{x} = \bar{X}(1 + k_1); \quad \bar{z} = \bar{Z}(1 + k_2);$$

with $E(k_0) = E(k_1) = E(k_2) = 0$

and

$$E(k_0^2) = \frac{\lambda S_y^2}{\bar{Y}^2}; \quad E(k_1^2) = \frac{\lambda S_x^2}{\bar{X}^2}; \quad E(k_2^2) = \frac{\lambda S_z^2}{\bar{Z}^2}; \quad E(k_0 k_1) = \frac{\lambda S_{yx}}{\bar{Y}\bar{X}}; \quad E(k_0 k_2) = \frac{\lambda S_{yz}}{\bar{Y}\bar{Z}};$$

$$E(k_1 k_2) = \frac{\lambda S_{zx}}{\bar{X}\bar{Z}},$$

Now expressing \bar{y}_{PR} in terms of k 's, we have

$$\bar{y}_{PR} = \bar{Y}(1 + k_0) \left[\theta(1 - gk_1)(1 - gk_2)^{-1} + (1 - \theta)(1 - gk_1)^{-1}(1 - gk_2) \right] \quad (2)$$

We assume that $|gk_1| < 1$ and $|gk_2| < 1$ so that the right hand side of (2) is expandable.

Now expanding the right hand side of (2) to the first degree of approximation, we have

$$\bar{y}_{PR} - \bar{Y} = \bar{Y} \left[k_0 + (1 - 2\alpha)g(k_1 - k_2 + k_0 k_1 - k_0 k_2) + g^2(k_1^2 - k_1 k_2 - \alpha(k_1^2 - k_2^2)) \right] \quad (3)$$

Taking expectations on both sides of (3), we get the bias of \bar{y}_{PR} to the first degree of approximation, as

$$B(\bar{y}_{PR}) = \lambda \bar{Y} \left[gDA + g^2(R_1^2 S_x^2 - R_1 R_2 S_{zx} - \theta(R_1^2 S_x^2 - R_2^2 S_z^2)) \right]$$

where $A = 1 - 2\theta$

Squaring both sides of (3) and neglecting terms of k 's involving power greater than two, we have

$$(\bar{y}_{PR} - \bar{Y})^2 = \bar{Y}^2 [k_0 + Agk_1 - Agk_2]^2$$

$$= \bar{Y}^2 [k_0^2 + 2Agk_0 k_1 - 2Agk_0 k_2 - 2A^2 g^2 k_1 k_2 + A^2 g^2 k_1^2 + A^2 g^2 k_2^2] \quad (4)$$

Taking expectations on both sides of (4), we get the MSE of \bar{y}_{PR} , to the first order of approximation, as

$$MSE(\bar{y}_{PR}) = \lambda[S_y^2 + 2AgD + A^2g^2C] \quad (5)$$

The MSE of the proposed estimator given in (5) can be re-written in terms of coefficient of variation as

$$MSE(\bar{y}_{PR}) = \lambda\bar{Y}^2[C_y^2 + 2AgC_yD^* + A^2g^2C^*]$$

where $C^* = C_x^2 + C_z^2 - 2\rho_{zx}C_zC_x$ and $D^* = \rho_{yx}C_x - \rho_{yz}C_z$, $C_y = \frac{S_y}{\bar{Y}}$, $C_x = \frac{S_x}{\bar{X}}$, $C_z = \frac{S_z}{\bar{Z}}$

The MSE equation given in (5) is minimized for

$$\theta = \frac{D + Cg}{2Cg} = \theta_0 \text{ (say)}$$

We can obtain the minimum MSE of the suggested estimator \bar{y}_{PR} , by using the optimal equation of θ in (5) as follows: $\min.MSE(\bar{y}_{PR}) = \lambda[S_y^2 + F(2D + CF)]$

where $F = g - E$ and $E = \frac{D + Cg}{C}$

3. EFFICIENCY COMPARISON

In this section, the efficiency of the suggested estimator \bar{y}_{PR} over the following estimator, \bar{y} , \bar{y}_R , \bar{y}_P , \bar{y}_S , \bar{y}_R^* , \bar{y}_P^* , \bar{y}_{SE} and \bar{y}_{ST} are investigated. We will have the conditions as follows:

$$(a) \quad MSE(\bar{y}_{PR}) - V(\bar{y}) < 0 \text{ if } \theta > \frac{2D + gC}{2gC}$$

$$(b) \quad MSE(\bar{y}_{PR}) - MSE(\bar{y}_R) < 0 \text{ if}$$

$$Ag(2D + AgC) < R_1(R_1S_x^2 - 2S_{yx}) \text{ provided } S_{yx} < \frac{R_1S_x^2}{2}$$

$$(c) \quad MSE(\bar{y}_{PR}) - MSE(\bar{y}_P) < 0 \text{ if}$$

$$Ag(2D + AgC) < R_2(R_2S_z^2 + 2S_{yz}) \text{ provided } S_{yz} < \frac{R_2S_z^2}{2}$$

$$(d) \text{ } MSE(\bar{y}_{PR}) - MSE(\bar{y}_S) < 0 \text{ if } C < \frac{-2D}{(Ag-1)} \text{ provided } g < \frac{1}{A}$$

$$(e) \text{ } MSE(\bar{y}_{PR}) - MSE(\bar{y}_R^*) < 0 \text{ if}$$

$$4\theta(\theta gC - gC - D) < 2gR_1R_2S_{zx} + 2R_2S_{yz} - 4R_1S_{yx} - gR_2^2S_z^2$$

$$(f) \text{ } MSE(\bar{y}_{PR}) - MSE(\bar{y}_P^*) < 0 \text{ if}$$

$$4\theta(\theta gC - gC - D) < 2gR_1R_2S_{zx} + 4R_2S_{yz} - 2R_1S_{yx} - gR_1^2S_x^2$$

$$(g) \text{ } MSE(\bar{y}_{PR}) - MSE(\bar{y}_{SE}) < 0 \text{ if } g < \frac{-2D}{C(A-1)} \text{ provided } A < 1$$

$$(h) \text{ } MSE(\bar{y}_{PR}) - MSE(\bar{y}_{ST}) < 0 \text{ if } Ag(2D + AgC) < -\rho^2S_y^2$$

4. NUMERICAL ILLUSTRATION

To analyze the performance of the suggested estimator in comparison to other estimators considered in this paper, two natural population data sets from the literature are being considered. The descriptions of these populations are given below.

(1) Population I [Singh (1969, p. 377)]; a detailed description can be seen in Singh (1965)

y : Number of females employed

x : Number of females in service

z : Number of educated females

$$N = 61, \quad n = 20, \quad \bar{Y} = 7.46, \quad \bar{X} = 5.31, \quad \bar{Z} = 179, \quad S_y^2 = 28.0818, \quad S_x^2 = 16.1761,$$

$$S_z^2 = 2028.1953, \quad \rho_{xy} = 0.7737, \quad \rho_{yz} = -0.2070, \quad \rho_{zx} = -0.0033,$$

(2) Population II [Source: Johnston 1972, p. 171]; A detailed description of these variables is shown in Table 1.

y : Percentage of hives affected by disease

x : Mean January temperature

z : Date of flowering of a particular summer species (number of days from January 1)

$N = 10, n = 4, \bar{Y} = 52, \bar{X} = 42, \bar{Z} = 200, S_y^2 = 65.9776, S_x^2 = 29.9880, S_z^2 = 84,$

$\rho_{xy} = 0.8, \rho_{yz} = -0.94, \rho_{zx} = -0.073,$

For these comparisons, the Percent Relative Efficiencies (PREs) of the different estimators are computed with respect to the usual unbiased estimator \bar{y} , using the formula

$$PRE(., \bar{y}) = \frac{V(\bar{y})}{MSE(.)} \times 100$$

and they are as presented in Table 2.

Table 1: Description of Population II.

y	x	z
49	35	200
40	35	212
41	38	211
46	40	212
52	40	203
59	42	194
53	44	194
61	46	188
55	50	196
64	50	190

Table 2 shows clearly that the proposed dual to ratio-cum-product estimator \bar{y}_{PR} has the highest PRE than other estimators; therefore, we can conclude based on the study populations that the suggested estimator is more efficient than the usual unbiased estimators, the traditional ratio and product estimator, ratio-cum-product estimator by Singh (1969),

Srivenkataramana (1980) estimator, Bandyopadhyay (1980) estimator, Singh et al. (2005) estimator and Sharma and Tailor (2010).

Table 2: PRE of the different estimators with respect to \bar{y}

Estimators	Population I	Population II
\bar{y}	100	100
\bar{y}_R	205	277
\bar{y}_P	102	187
\bar{y}_S	214	395
\bar{y}_R^*	215	239
\bar{y}_P^*	105	150
\bar{y}_{SE}	236	402
\bar{y}_{ST}	250	278
\bar{y}_{PR}	279	457

5. CONCLUSION

We have developed a new estimator for estimating the finite population mean, which is found to be more efficient than the usual unbiased estimator, the traditional ratio and product estimators and the estimators proposed by Singh (1969), Srivenkataramana (1980), Bandyopadhyay (1980), Singh et al. (2005) and Sharma and Tailor (2010). This theoretical inference is also satisfied by the result of an application with original data. In future, we hope to extend the estimators suggested here for the development of a new estimator in stratified random sampling.

REFERENCES

1. ABU-DAYYEH, W. A., AHMED, M. S., AHMED, R. A., and MUTTLAK, H. A. (2003): Some estimators of finite population mean using auxiliary information, *Applied Mathematics and Computation*, 139, 287–298.
2. BANDYOPADHYAY, S. (1980): Improved ratio and product estimators, *Sankhya series C* 42, 45- 49.
3. DIANA, G. and PERRI, P.F. (2007): Estimation of finite population mean using multi-auxiliary information. *Metron*, Vol. LXV, Number 1, 99-112
4. JOHNSTON, J. (1972): *Econometric methods*, (2nd edn), McGraw-Hill, Tokyo.
5. KADILAR, C. and CINGI, H. (2004): Estimator of a population mean using two auxiliary variables in simple random sampling, *International Mathematical Journal*, 5, 357–367.
6. KADILAR, C. and CINGI, H. (2005): A new estimator using two auxiliary variables, *Applied Mathematics and Computation*, 162, 901–908.
7. Malik, S. and Singh, R. (2012) : Some improved multivariate ratio type estimators using geometric and harmonic means in stratified random sampling. *ISRN Prob. and Stat.* Article ID 509186, doi:10.5402/2012/509186.
8. MURTHY, M. N. (1964): Product method of estimation. *Sankhya*, A, 26, 294-307.
9. OLKIN, I. (1958): Multivariate ratio estimation for finite populations, *Biometrika*, 45, 154–165.
10. PERRI, P. F. (2004): Alcune considerazioni sull’efficienza degli stimatori rapporto-cum-prodotto, *Statistica & applicazioni*, 2, no. 2, 59–75.
11. PERRI, P. F. (2005): Combining two auxiliary variables in ratio-cum-product type estimators. *Proceedings of Italian Statistical Society, Intermediate Meeting on Statistics and Environment*, Messina, 21-23 September 2005, 193–196.
12. RAJ, D. (1965): On a method of using multi-auxiliary information in sample surveys, *Journal of the American Statistical Association*, 60, 154–165.
13. REDDY, V. N. (1974): On a transformed ratio method of estimation, *Sankhya C*, **36**, 59–70.
14. SHARMA, B. and TAILOR, R. (2010): A New Ratio-Cum-Dual to Ratio Estimator of Finite Population Mean in Simple Random Sampling. *Global Journal of Science Frontier Research*, Vol. 10, Issue 1, 27-31
15. SINGH, M.P. (1965): On the estimation of ratio and product of the population parameters, *Sankhya series B* 27, 321-328.
16. SINGH, M. P. (1967): Multivariate product method of estimation for finite populations, *Journal of the Indian Society of Agricultural Statistics*, 31, 375–378.
17. SINGH, M.P. (1969): Comparison of some ratio-cum-product estimators, *Sankhya series B* 31, 375-378.

18. SINGH, H.P., SINGH, R., ESPEJO, M.R., PINEDA, M.D and NADARAJAH, S. (2005): On the efficiency of a dual to ratio-cum-product estimator in sample surveys. *Mathematical Proceedings of the Royal Irish Academy*, 105A (2), 51-56.
19. SINGH, R., SINGH, H.P AND ESPEJO, M.R. (1998): The efficiency of an alternative to ratio estimator under a super population model. *J.S.P.I.*, 71, 287-301.
20. SINGH, R., CHAUHAN, P. AND SAWAN, N. (2007): On the bias reduction in linear variety of alternative to ratio-cum-product estimator. *Statistics in Transition*,8(2),293-300.
21. SINGH, R., KUMAR, M., CHAUHAN, P., SAWAN, N. AND SMARANDACHE, F. (2011): A general family of dual to ratio-cum-product estimator in sample surveys. *SIT_newseries*, 12(3), 587-594.
22. SRIVENKATARAMANA, T. (1978): Change of origin and scale in ratio and difference methods of estimation in sampling, *Canad. J. Stat.*, **6**, 79–86.
23. SRIVENKATARAMANA, T. (1980): A dual to ratio estimator in sample surveys, *Biometrika*, **67**(1), 199–204.
24. TRACY, D.S., SINGH, H.P. AND SINGH, R. (1996): An alternative to ratio-cum-product estimator in sample surveys. *J.S.P.I.*,53,375-387.

Some Improved Estimators for Population Variance Using Two Auxiliary Variables in Double Sampling

¹Viplav Kumar Singh, †¹Rajesh Singh and ²Florentin Smarandache

¹Department of Statistics, Banaras Hindu University, Varanasi-221005, India

²Department of Mathematics, University of New Mexico, Gallup, USA

†*Corresponding author*

Abstract

In this article we have proposed an efficient generalised class of estimator using two auxiliary variables for estimating unknown population variance S_y^2 of study variable y . We have also extended our problem to the case of two phase sampling. In support of theoretical results we have included an empirical study.

1. Introduction

Use of auxiliary information improves the precision of the estimate of parameter. Out of many ratio and product methods of estimation are good example in this context. We can use ratio method of estimation when correlation coefficient between auxiliary and study variate is positive (high), on the other hand we use product method of estimation when correlation coefficient between auxiliary and study variate is highly negative.

Variations are present everywhere in our day-to-day life. An agriculturist needs an adequate understanding of the variations in climatic factors especially from place to place (or time to time) to be able to plan on when, how and where to plant his crop. The problem of estimation of finite population variance S_y^2 , of the study variable y was discussed by Isaki (1983), Singh and Singh (2001, 2002, 2003), Singh et al. (2008), Grover (2010), and Singh et al. (2011).

Let x and z are auxiliary variates having values (x_i, z_i) and y is the study variate having values (y_i) respectively. Let $V_i (i = 1, 2, \dots, N)$ is the population having N units such that y is positively correlated x and negatively correlated with z . To estimate S_y^2 , we assume that S_x^2 and S_z^2 are known, where

$$S_y^2 = \frac{1}{N} \sum_{i=1}^N (y_i - \bar{Y})^2, \quad S_x^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \bar{X})^2 \text{ and } S_z^2 = \frac{1}{N} \sum_{i=1}^N (z_i - \bar{Z})^2.$$

Assume that N is large so that the finite population correction terms are ignored. A sample of size n is drawn from the population V using simple random sample without replacement.

Usual unbiased estimator of population variance S_y^2 is s_y^2 , where, $s_y^2 = \frac{1}{(n-1)} \sum_{i=1}^n (y_i - \bar{y})^2$.

Up to the first order of approximation, variance of s_y^2 is given by

$$\text{var}(s_y^2) = \frac{S_y^4}{n} \partial_{400}^* \quad (1.1)$$

where, $\partial_{400}^* = \partial_{400} - 1$, $\partial_{pqr} = \frac{\mu_{pqr}}{\mu_{200}^{p/2} \mu_{020}^{q/2} \mu_{002}^{r/2}}$, and

$\mu_{pqr} = \frac{1}{N} \sum_{i=1}^N (y_i - \bar{Y})^p (x_i - \bar{X})^q (z_i - \bar{Z})^r$; p, q, r being the non-negative integers.

2. Existing Estimators

Let $s_y^2 = S_y^2(1 + e_0)$, $s_x^2 = S_x^2(1 + e_1)$ and $s_z^2 = S_z^2(1 + e_2)$

where, $s_x^2 = \frac{1}{(n-1)} \sum_{i=1}^n (x_i - \bar{x})^2$, $s_z^2 = \frac{1}{(n-1)} \sum_{i=1}^n (z_i - \bar{z})^2$

and $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$, $\bar{z} = \frac{1}{n} \sum_{i=1}^n z_i$.

Also, let

$$E(e_1) = E(e_2) = 0$$

$$E(e_0^2) = \frac{\partial_{400}^*}{n}, E(e_1^2) = \frac{\partial_{040}^*}{n} \text{ and } E(e_2^2) = \frac{\partial_{004}^*}{n}$$

$$E(e_0 e_1) = \frac{1}{n} \partial_{220}^*, E(e_1 e_2) = \frac{1}{n} \partial_{022}^*, E(e_0 e_2) = \frac{1}{n} \partial_{202}^*$$

Isaki (1983) suggested ratio estimator t_1 for estimating S_y^2 as-

$$t_1 = s_y^2 \frac{S_y^2}{S_x^2}; \text{ where } s_x^2 \text{ is unbiased estimator of } S_x^2 \quad (1.2)$$

Up to the first order of approximation, mean square error of t_1 is given by,

$$MSE(t_1) = \frac{S_y^4}{n} [\partial_{400}^* + \partial_{040}^* - 2\partial_{220}^*] \quad (1.3)$$

Singh et al. (2007) proposed the exponential ratio-type estimator t_2 as-

$$t_2 = s_y^2 \exp \left[\frac{S_x^2 - s_x^2}{S_x^2 + s_x^2} \right] \quad (1.4)$$

And exponential product type estimator t_3 as-

$$t_3 = s_y^2 \left[\frac{s_x^2 - S_x^2}{s_x^2 + S_x^2} \right] \quad (1.5)$$

Following Kadilar and Cingi (2006), Singh et al. (2011) proposed an improved estimator for estimating population variance S_y^2 , as-

$$t_4 = s_y^2 \left[k_4 \exp \left\{ \frac{S_x^2 - s_x^2}{S_x^2 + s_x^2} \right\} + (1 - k_4) \left\{ \frac{s_x^2 - S_x^2}{s_x^2 + S_x^2} \right\} \right] \quad (1.6)$$

where k_4 is a constant.

Up to the first order of approximation mean square errors of t_2, t_3 and t_4 are respectively given by

$$MSE(t_2) = \frac{S_y^4}{n} \left[\partial_{400}^* + \frac{\partial_{040}^*}{4} - \partial_{220}^* \right] \quad (1.7)$$

$$\text{MSE}(t_3) = \frac{S_y^4}{n} \left[\partial_{400}^* + \frac{\partial_{004}^*}{4} - \partial_{202}^* \right] \quad (1.8)$$

$$\text{MSE}(t_4) = \frac{S_y^4}{n} \left[\partial_{400}^* + k_4^2 \frac{\partial_{040}^*}{4} + (1-k_4)^2 \frac{\partial_{004}^*}{4} - k_4 \partial_{220}^* + (1-k_4) \partial_{202}^* - \frac{k_4(1-k_4)}{2} \partial_{022}^* \right] \quad (1.9)$$

$$\text{where } k_4 = \frac{(\partial_{004}^*/2) + \partial_{220}^* + \partial_{022}^*}{2(\partial_{040}^* + \partial_{004}^* + \partial_{022}^*)}.$$

3. Improved Estimator

Using Singh and Solanki (2011), we propose some improved estimators for estimating population variance S_y^2 as-

$$t_5 = s_y^2 \left[\frac{cS_x^2 - Ds_x^2}{(c-d)S_x^2} \right]^p \quad (1.10)$$

$$t_6 = s_y^2 \left[\frac{(a+b)S_z^2}{aS_x^2 + bS_x^2} \right]^q \quad (1.11)$$

$$t_7 = s_y^2 \left[k_7 \left\{ \frac{cS_x^2 - Ds_x^2}{(c-d)S_x^2} \right\}^p + (1-k_7) \left\{ \frac{(a+b)S_z^2}{aS_x^2 + bS_x^2} \right\}^q \right] \quad (1.12)$$

where a, b, c, d are suitably chosen constants and k_7 is a real constant to be determined so as to minimize MSE's.

Expressing t_5, t_6 and t_7 in terms of e_i 's, we have

$$t_5 = S_y^2 [1 - x_1 p e_1 + e_0 - x_1 p e_0 e_1] \quad (1.13)$$

$$\text{where, } x_1 = \frac{d}{(c-d)}.$$

$$t_6 = S_y^2 [1 - q x_2 e_2 + e_0 - q x_2 e_0 e_2] \quad (1.14)$$

$$\text{where, } x_2 = \frac{b}{(a+b)}.$$

$$t_7 = S_y^2 [1 + e_0 + p x_1 k_7 (e_1' - e_1) + (k_7 - 1) q x_2 e_2'] \quad (1.15)$$

The mean squared error of estimators are obtained by subtracting S_y^2 from each estimator and squaring both sides and than taking expectations-

$$\text{MSE}(t_5) = \frac{S_y^4}{n} [\partial_{400}^* + x_1^2 p^2 \partial_{040}^* - 2x_1 p \partial_{220}^*] \quad (1.16)$$

Differentiating (1.16) with respect to x_1 , we get the optimum value of x_1 as-

$$x_{1(\text{opt})} = \frac{\partial_{220}^*}{p \partial_{040}^*}.$$

$$\text{MSE}(t_6) = \frac{S_y^4}{n} [\partial_{400}^* + x_2^2 q^2 \partial_{004}^* - 2x_2 q \partial_{202}^*] \quad (1.17)$$

Differentiating (1.17) with respect to x_2 , we get the optimum value of x_2 as –

$$x_{2(\text{opt})} = \frac{\partial_{202}^*}{q \partial_{004}^*}.$$

$$\text{MSE}(t_7) = \frac{S_y^4}{n} [A + k_7^2 B + (1 - k_7)C - 2k_7(1 - k_7)E - 2(1 - k_7)F] \quad (1.18)$$

Differentiating (1.18) with respect to k_7 , we get the optimum value k_7 of as –

$$k_{7(\text{opt})} = \frac{C + D - F - E}{B + C - 2E}.$$

where

$$A = \partial_{400}^*, \quad B = x_1^2 p^2 \partial_{040}^*,$$

$$C = x_2^2 q^2 \partial_{004}^*, \quad D = x_2 q \partial_{202}^*,$$

$$E = x_1 x_2 p q \partial_{022}^*, \quad F = x_2 q \partial_{202}^*.$$

2. Estimators In Two Phase Sampling

In certain practical situations when S_x^2 is not known a priori, the technique of two phase sampling or double sampling is used. Allowing SRSWOR design in each phase, the two – phase sampling scheme is as follows:

- The first phase sample s'_n ($s'_n \subset V$) of a fixed size n' is drawn to measure only x and z in order to formulate the a good estimate of S_x^2 and S_z^2 , respectively.

- Given s'_n , the second phase sample s_n ($s_n \subset s'_n$) of a fixed size n is drawn to measure y only.

Existing Estimators

Singh et al. (2007) proposed some estimators to estimate S_y^2 in two phase sampling, as:

$$t'_2 = s_y^2 \exp \left[\frac{s_x'^2 - s_x^2}{s_x'^2 + s_x^2} \right] \quad (2.1)$$

$$t'_3 = s_y^2 \exp \left[\frac{s_z'^2 - s_z^2}{s_z'^2 + s_z^2} \right] \quad (2.2)$$

$$t'_4 = s_y^2 \left[k'_4 \exp \left\{ \frac{s_x'^2 - s_x^2}{s_x'^2 + s_x^2} \right\} + (1 - k'_4) \exp \left\{ \frac{s_z'^2 - s_z^2}{s_z'^2 + s_z^2} \right\} \right] \quad (2.3)$$

MSE of the estimator t'_2 , t'_3 and t'_4 are respectively, given by

$$\text{MSE}(t'_2) = S_y^4 \left[\frac{\partial_{400}^*}{n} + \frac{1}{4} \left(\frac{1}{n} - \frac{1}{n'} \right) \partial_{040}^* + \left(\frac{1}{n'} - \frac{1}{n} \right) \partial_{220}^* \right] \quad (2.4)$$

$$\text{MSE}(t'_3) = S_y^4 \left[\frac{\partial_{400}^*}{n} + \frac{1}{4} \left(\frac{1}{n} - \frac{1}{n'} \right) \partial_{004}^* - \left(\frac{1}{n'} - \frac{1}{n} \right) \partial_{202}^* \right] \quad (2.5)$$

$$\text{MSE}(t'_4) = S_y^4 [A' + k_4'^2 B' + (1 - k_4')^2 C' + k_4' D' + (1 - k_4') E'] \quad (2.6)$$

$$\text{And } k_{4(\text{opt})}' = \frac{2C' + E' - D'}{2(B' + C')}$$

Where,

$$A' = \frac{\partial_{400}^*}{n}, B' = \frac{1}{4} \left(\frac{1}{n} - \frac{1}{n'} \right) \partial_{040}^*, C' = \frac{1}{4n'} \partial_{004}^*$$

$$D' = \left(\frac{1}{n'} - \frac{1}{n} \right) \partial_{202}^*, E' = \frac{1}{n'} \partial_{202}^*$$

Proposed estimators in two phase sampling

The estimator proposed in section 3 will take the following form in two phase sampling:

$$t'_5 = s_y^2 \left[\frac{cs_x'^2 - ds_x^2}{(c-d)s_x'^2} \right] \quad (2.7)$$

$$t'_6 = S_y^2 \left[\frac{(a+b)S_z^2}{aS_z^2 + bs_z'^2} \right] \quad (2.8)$$

$$t'_7 = S_y^2 \left[k'_7 \left\{ \frac{cs_x'^2 - ds_x^2}{(c-d)s_x'^2} \right\} + (1-k'_7) \left\{ \frac{(a+b)S_z^2}{aS_z^2 + bs_z'^2} \right\} \right] \quad (2.9)$$

Let,

$$s_y^2 = S_y^2(1+e_0), s_x^2 = S_x^2(1+e_1), s_x'^2 = S_x^2(1+e_1')$$

$$s_z^2 = S_z^2(1+e_2), s_z'^2 = S_z^2(1+e_2')$$

Where,

$$s_x'^2 = \frac{1}{(n'-1)} \sum_{i=1}^{n'} (x_i - \bar{x}')^2, s_z'^2 = \frac{1}{(n'-1)} \sum_{i=1}^{n'} (z_i - \bar{z}')^2$$

$$\text{and } \bar{x}' = \frac{1}{n'} \sum_{i=1}^{n'} x_i, \bar{z}' = \frac{1}{n'} \sum_{i=1}^{n'} z_i$$

Also,

$$E(e_1') = E(e_2') = 0$$

$$E(e_1'^2) = \frac{\partial_{040}^*}{n'}, E(e_2'^2) = \frac{\partial_{004}^*}{n'}$$

$$E(e_0 e_1') = \frac{1}{n'} \partial_{220}^*, E(e_0 e_2') = \frac{1}{n'} \partial_{202}^*, E(e_1 e_1') = \frac{1}{n'} \partial_{040}^*$$

$$E(e_2 e_2') = \frac{1}{n'} \partial_{004}^*, E(e_1 e_2') = \frac{1}{n'} \partial_{022}^*, E(e_1' e_2') = \frac{1}{n'} \partial_{022}^*$$

Writing estimators t'_5, t'_6 and t'_7 in terms of e_i 's we have ,respectively

$$t'_5 = S_y^2 [1 + e_0 + px_1(e_1' - e_1)] \quad (2.10)$$

$$t'_6 = S_y^2 [1 + e_0 - qx_2 e_2'] \quad (2.11)$$

$$t'_7 = S_y^2 [1 + px_1 k'_7 ((e_1' - e_1) + e_0 + (k'_7 - 1)qx_2 e_2')] \quad (2.12)$$

Solving (1.10),(1.11) and (1.12),we get the MSE'S of the estimators t'_5, t'_6 and t'_7 , respectively as-

$$MSE(t'_5) = S_y^4 \left[\frac{\partial_{400}^*}{n} + p^2 x_1^2 \left(\frac{1}{n} - \frac{1}{n'} \right) \partial_{040}^* + 2px_1 \left(\frac{1}{n'} - \frac{1}{n} \right) \partial_{220}^* \right] \quad (2.13)$$

Differentiate (2.13) w.r.t. x_1 , we get the optimum value of x_1 as-

$$x_{1(\text{opt})} = \frac{\partial_{220}^*}{p\partial_{040}^*}$$

$$\text{MSE}(t'_6) = S_y^4 \left[\frac{\partial_{400}^*}{n} + q^2 x_2^2 \frac{1}{n'} \partial_{004}^* + 2qx_2 \frac{1}{n'} \partial_{202}^* \right] \quad (2.14)$$

Differentiate (2.14) with respect to x_2 , we get the optimum value of x_2 as –

$$x_{2(\text{opt})} = \frac{\partial_{202}^*}{q\partial_{004}^*}.$$

$$\text{MSE}(t'_7) = S_y^4 \left[A_1 + B_1 k_7'^2 + (k_7' - 1)^2 C_1 + 2k_7' D_1 + 2(k_7' - 1) E_1 \right] \quad (2.15)$$

Differentiate (2.15) with respect to k_7' , we get the optimum value of k_7' as –

$$k_{7(\text{opt})}' = \frac{C_1 - D_1 - E_1}{B_1 + C_1}$$

Where,

$$A_1 = \frac{\partial_{400}^*}{n}, B_1 = p^2 x_1^2 \left(\frac{1}{n} - \frac{1}{n'} \right) \partial_{040}^*, C_1 = px_1 \frac{q^2 x_2^2}{n'} \partial_{004}^*$$

$$D_1 = px_1 \left(\frac{1}{n'} - \frac{1}{n} \right) \partial_{220}^*, E_1 = qx_2 \frac{\partial_{202}^*}{n'}$$

5. Empirical Study

In support of theoretical result an empirical study is carried out. The data is taken from Murthy(1967):

$$\partial_{400} = 3.726, \partial_{040} = 2.912, \partial_{044} = 2.808$$

$$\partial_{022} = 2.73, \partial_{202} = 2.979, \partial_{220} = 3.105$$

$$c_x = 0.5938, c_y = 0.7531, c_z = 0.7205$$

$$\rho_{yz} = 0.904, \rho_{xy} = 0.98, n = 7, n' = 15$$

$$\bar{X} = 747.5882, \bar{Y} = 199.4412, \bar{Z} = 208.8824$$

❖ In Table 5.1 percent relative efficiency of various estimators of S_y^2 is written with respect to s_y^2

➤ **Table 5.1: PRE of the estimator with respect to s_y^2**

Estimators	PRE
S_y^2	100
t_1	636.9158
t_2	248.0436
t_3	52.86019
t_4	699.2526
t_5	667.2895
t_6	486.9362
t_7	699.5512

- ❖ In Table 5.2 percent relative efficiency of various estimators of S_y^2 is written with respect to s_y^2 in two phase sampling:

➤ **Table 5.2: PRE of the estimators in two phase sampling with respect to S_y^2**

Estimators	PRE
S_y^2	100
t_2'	142.60
t_3'	66.42
t_4'	460.75
t_5'	182.95
t_6'	158.93
t_7'	568.75

6. Conclusion

In Table 5.1 and 5.2 percent relative efficiencies of various estimators are written with respect to s_y^2 . From Table 5.1 and 5.2 we observe that the proposed estimator

under optimum condition performs better than usual estimator, Isaki (1983) estimator and Singh et al. (2007) estimator.

7. References

Bahl, S. and Tuteja, R.K. (1991): Ratio and Product type exponential estimator. *Information and Optimization sciences*, Vol. XII, I, 159-163.

Grover, L. K . (2010): A correction note on improvement in variance estimation using auxiliary information. *Commun. Stat. Theo. Meth.* 39:753–764.

Isaki, C. T. (1983): Variance estimation using auxiliary information. *Journal of American Statistical Association*.

Kadilar, C. and Cingi, H. (2006) : Improvement in estimating the population mean in simple random sampling. *Applied Mathematics Letters* 19 (2006), 75–79.

Murthy, M. N. (1967):. *Sampling Theory and Methods*, Statistical Publishing Soc., Calcutta, India.

Singh, H. P. and Singh, R. (2001): Improved Ratio-type Estimator for Variance Using Auxiliary Information. *J.I.S.A.S.*,54(3),276-287.

Singh, H. P. and Singh, R. (2002): A class of chain ratio-type estimators for the coefficient of variation of finite population in two phase sampling. *Aligarh Journal of Statistics*, 22,1-9.

Singh, H. P. and Singh, R. (2003): Estimation of variance through regression approach in two phase sampling. *Aligarh Journal of Statistics*, 23, 13-30.

Singh, H. P, and Solanki, R. S. (2012): A new procedure for variance estimation in simple random sampling using auxiliary information. *Statistical papers*, DOI 10.1007/s00362-012-0445-2.

Singh, R., Chauhan, P., Sawan, N. and Smarandache, F. (2011): Improved exponential estimator for population variance using two auxiliary variables. *Italian Jour. Of Pure and Applied. Math.*, 28, 103-110.

Singh, R., Kumar, M., Singh, A.K. and Smarandache, F. (2011): A family of estimators of population variance using information on auxiliary attribute. *Studies in sampling techniques and time series analysis*. Zip publishing, USA.

Singh, R., Chauhan, P., Sawan, N. and Smarandache, F. (2008): Almost unbiased ratio and product type estimator of finite population variance using the knowledge of kurtosis of an auxiliary variable in sample surveys. *Octagon Mathematical Journal*, Vol. 16, No. 1, 123-130.

The purpose of writing this book is to suggest some improved estimators using auxiliary information in sampling schemes like simple random sampling and systematic sampling.

This volume is a collection of five papers, written by eight coauthors (listed in the order of the papers): Manoj K. Chaudhary, Sachin Malik, Rajesh Singh, Florentin Smarandache, Hemant Verma, Prayas Sharma, Olufadi Yunusa, and Viplav Kumar Singh, from India, Nigeria, and USA.

The following problems have been discussed in the book:

In chapter one an estimator in systematic sampling using auxiliary information is studied in the presence of non-response. In second chapter some improved estimators are suggested using auxiliary information. In third chapter some improved ratio-type estimators are suggested and their properties are studied under second order of approximation.

In chapter four and five some estimators are proposed for estimating unknown population parameter(s) and their properties are studied.

This book will be helpful for the researchers and students who are working in the field of finite population estimation.

