

# Neutrosophic Crisp Sets & Neutrosophic Crisp Topological Spaces

A. A. Salama<sup>1</sup>, Florentin Smarandache<sup>2</sup> and Valeri Kroumov<sup>3</sup>

<sup>1</sup> Department of Mathematics and Computer Science, Faculty of Sciences,  
Port Said University, Egypt, e-mail: drsalama44@gmail.com

<sup>2</sup> Department of Mathematics, University of New Mexico Gallup, NM, USA,  
e-mail: smarand@unm.edu

<sup>3</sup> Department of Electrical and Electronic Engineering,  
Okayama University of Science, Okayama, Japan, e-mail: val@ee.ous.ac.jp

December 27, 2013

**Abstract.** In this paper, we generalize the crisp topological space to the notion of neutrosophic crisp topological space, and we construct the basic concepts of the neutrosophic crisp topology. In addition to these, we introduce the definitions of neutrosophic crisp continuous function and neutrosophic crisp compact spaces. Finally, some characterizations concerning neutrosophic crisp compact spaces are presented and one obtains several properties. Possible application to GIS topology rules are touched upon.

**Keywords:** Neutrosophic Crisp Set; Neutrosophic Topology; Neutrosophic Crisp Topology.

## 1 Introduction

Neutrosophy has laid the foundation for a whole family of new mathematical theories generalizing both their crisp and fuzzy counterparts, such as a neutrosophic set theory in [1, 2, 3]. After the introduction of the neutrosophic set concepts in [4, 5, 6, 7, 8, 9, 10, 11, 12] and later have given the fundamental definitions of neutrosophic set operations we generalize the crisp topological space to the notion of neutrosophic crisp set. Finally, we introduce the definitions of neutrosophic crisp continuous function and neutrosophic crisp compact space, and we obtain several properties and some characterizations concerning the neutrosophic crisp compact space.

## 2 Terminologies

We recollect some relevant basic preliminaries, and in particular, the work of Smarandache in [1, 2, 3, 11], and Salama et al. [4, 5, 6, 7, 8, 11]. Smarandache introduced the neutrosophic components T, I, F which represent the membership, indeterminacy, and non-membership values respectively, where  $]^{-0, 1^{+}[$  is a non-standard unit interval.

Hanafy and Salama *et al.*[10, 11] considered some possible definitions for basic concepts of the neutrosophic crisp set and its operations. We now improve some results by the following.

## 3 Neutrosophic Crisp Sets

**Definition 3.1** *Let  $X$  be a non-empty fixed set. A neutrosophic crisp set (NCS)  $A$  is an object having the form  $A = \langle A_1, A_2, A_3 \rangle$ , where  $A_1, A_2$ , and  $A_3$  are subsets of  $X$  satisfying  $A_1 \cap A_2 = \phi$ ,  $A_1 \cap A_3 = \phi$ , and  $A_2 \cap A_3 = \phi$ .*

**Remark 3.1** *A neutrosophic crisp set  $A = \langle A_1, A_2, A_3 \rangle$  can be identified as an ordered triple  $\langle A_1, A_2, A_3 \rangle$ , where  $A_1, A_2$ , and  $A_3$  are subsets on  $X$ , and one can define several relations and operations between NCSs.*

Since our purpose is to construct the tools for developing neutrosophic crisp sets, we must introduce the types of NCSs  $\phi_N$  and  $X_N$  in  $X$  as follows:

1)  $\phi_N$  may be defined in many ways as an NCS, as follows

- i)  $\phi_N = \langle \phi, \phi, X \rangle$ , or
- ii)  $\phi_N = \langle \phi, X, X \rangle$ , or
- iii)  $\phi_N = \langle \phi, X, \phi \rangle$ , or
- iv)  $\phi_N = \langle \phi, \phi, \phi \rangle$ .

2)  $X_N$  may also be defined in many ways as an NCS:

- i)  $X_N = \langle X, \phi, \phi \rangle$ ,
- ii)  $X_N = \langle X, X, \phi \rangle$ ,
- iii)  $X_N = \langle X, X, X \rangle$ .

Every crisp set  $A$  formed by three disjoint subsets of a non-empty set  $X$  is obviously an NCS having the form  $A = \langle A_1, A_2, A_3 \rangle$ .

**Definition 3.2** Let  $A = \langle A_1, A_2, A_3 \rangle$  an NCS on  $X$ , then the complement  $A^c$  of the set  $A$  may be defined in three different ways:

$$(C_1) A^c = \langle A_1^c, A_2^c, A_3^c \rangle,$$

$$(C_2) A^c = \langle A_3, A_2, A_1 \rangle,$$

$$(C_3) A^c = \langle A_3, A_2^c, A_1 \rangle.$$

One can define several relations and operations between NCSs as follows:

**Definition 3.3** Let  $X$  be a non-empty set, and NCSs  $A$  and  $B$  in the form  $A = \langle A_1, A_2, A_3 \rangle$ ,  $B = \langle B_1, B_2, B_3 \rangle$ , then we may consider two possible definitions for subsets ( $A \subseteq B$ ):

- 1)  $A \subseteq B \Leftrightarrow A_1 \subseteq B_1, A_2 \subseteq B_2$  and  $A_3 \supseteq B_3$ ,  
or
- 2)  $A \subseteq B \Leftrightarrow A_1 \subseteq B_1, A_2 \supseteq B_2$  and  $A_3 \supseteq B_3$ .

**Proposition 3.1** For any neutrosophic crisp set  $A$  the following are hold:

- i)  $\phi_N \subseteq A, \phi_N \subseteq \phi_N$ ,
- ii)  $A \subseteq X_N, X_N \subseteq X_N$ .

**Definition 3.4** Let  $X$  is a non-empty set, and the NCSs  $A$  and  $B$  in the form  $A = \langle A_1, A_2, A_3 \rangle$ ,  $B = \langle B_1, B_2, B_3 \rangle$ . Then:

- 1)  $A \cap B$  may be defined in two ways:
  - i)  $A \cap B = \langle A_1 \cap B_1, A_2 \cap B_2, A_3 \cup B_3 \rangle$  or
  - ii)  $A \cap B = \langle A_1 \cap B_1, A_2 \cup B_2, A_3 \cup B_3 \rangle$ .
- 2)  $A \cup B$  may also be defined in two ways:
  - i)  $A \cup B = \langle A_1 \cup B_1, A_2 \cap B_2, A_3 \cap B_3 \rangle$  or
  - ii)  $A \cup B = \langle A_1 \cup B_1, A_2 \cup B_2, A_3 \cap B_3 \rangle$ .
- 3)  $[ ]A = \langle A_1, A_2, A_1^c \rangle$ .
- 4)  $\langle \rangle A = \langle A_3^c, A_2, A_3 \rangle$ .

**Proposition 3.2** For any two neutrosophic crisp sets  $A$  and  $B$  on  $X$ , the followings are true:

- 1)  $(A \cap B)^c = A^c \cup B^c$ .
- 2)  $(A \cup B)^c = A^c \cap B^c$ .

We can easily generalize the operations of intersection and union in Definition 3.2 to arbitrary family of neutrosophic crisp subsets as follows:

**Proposition 3.3** *Let  $\{A_j : j \in J\}$  be arbitrary family of neutrosophic crisp subsets in  $X$ . Then*

1)  $\cap A_j$  may be defined as the following types:

- i)  $\cap A_j = \langle \cap A_{J1}, \cap A_{J2}, \cup A_{j3} \rangle$ , or
- ii)  $\cap A_j = \langle \cap A_{J1}, \cup A_{J2}, \cup A_{j3} \rangle$ .

2)  $\cup A_j$  may be defined as the following types:

- i)  $\cup A_j = \langle \cup A_{J1}, \cap A_{J2}, \cap A_{j3} \rangle$ , or
- ii)  $\cup A_j = \langle \cup A_{J1}, \cup A_{J2}, \cap A_{j3} \rangle$ .

**Definition 3.5** *The product of two neutrosophic crisp sets  $A$  and  $B$  is a neutrosophic crisp set given by*

$$A \times B = \langle A_1 \times B_1, A_2 \times B_2, A_3 \times B_3 \rangle$$

## 4 Neutrosophic crisp Topological Spaces

Here we extend the concepts of topological space and intuitionistic topological space to the case of neutrosophic crisp sets.

**Definition 4.1** *A neutrosophic crisp topology (NCT) on a non-empty set  $X$  is a family  $\Gamma$  of neutrosophic crisp subsets in  $X$  satisfying the following axioms*

- i)  $\phi_N, X_N \in \Gamma$ .
- ii)  $A_1 \cap A_2 \in \Gamma$  for any  $A_1$  and  $A_2 \in \Gamma$ .
- iii)  $\cup A_j \in \Gamma \quad \forall \{A_j : j \in J\} \subseteq \Gamma$ .

In this case the pair  $(X, \Gamma)$  is called a neutrosophic crisp topological space (NCTS) in  $X$ . The elements in  $\Gamma$  are called neutrosophic crisp open sets (NCOSs) in  $Y$ . A neutrosophic crisp set  $F$  is closed if and only if its complement  $F^c$  is an open neutrosophic crisp set.

**Remark 4.1** *Neutrosophic crisp topological spaces are very natural generalizations of topological spaces and intuitionistic topological spaces, and they allow more general functions to be members of topology:*

$$TS \rightarrow ITS \rightarrow NCTS$$

**Example 4.1** *Let  $X = \{a, b, c, d\}$ ,  $\phi_N, X_N$  be any types of the universal and empty subsets, and  $A, B$  two neutrosophic crisp subsets on  $X$  defined by  $A = \langle \{a\}, \{b, d\}, c \rangle$ ,  $B = \langle \{a\}, \{b\}, \{c\} \rangle$ , then the family  $\Gamma = \{\phi_N, X_N, A, B\}$  is a neutrosophic crisp topology on  $X$ .*

**Example 4.2** *Let  $(X, \tau_0)$  be a topological space such that  $\tau_0$  is not indiscrete. Suppose  $\{G_i : i \in J\}$  be a family and  $\tau_0 = \{X, \phi\} \cup \{G_i : i \in J\}$ . Then we can construct the following topologies:*

i) *Two intuitionistic topologies*

- a)  $\tau_1 = \{\phi_I, X_I\} \cup \{\langle G_i, \phi \rangle, i \in J\}$ .
- b)  $\tau_2 = \{\phi_I, X_I\} \cup \{\langle \phi, G_i^c \rangle, i \in J\}$ .

ii) *Four neutrosophic crisp topologies*

- a)  $\Gamma_1 = \{\phi_N, X_N\} \cup \{\langle \phi, \phi, G_i^c \rangle, i \in J\}$ .
- b)  $\Gamma_2 = \{\phi_N, X_N\} \cup \{\langle G_i, \phi, \phi \rangle, i \in J\}$ .
- c)  $\Gamma_3 = \{\phi_N, X_N\} \cup \{\langle G_i, \phi, G_i^c \rangle, i \in J\}$ .
- d)  $\Gamma_4 = \{\phi_N, X_N\} \cup \{\langle G_i^c, \phi, \phi \rangle, i \in J\}$ .

**Definition 4.2** Let  $(X, \Gamma_1), (X, \Gamma_2)$  be two neutrosophic crisp topological spaces on  $X$ . Then  $\Gamma_1$  is said to be contained in  $\Gamma_2$  (in symbols  $\Gamma_1 \subseteq \Gamma_2$ ) if  $G \in \Gamma_2$  for each  $G \in \Gamma_1$ . In this case, we also say that  $\Gamma_1$  is coarser than  $\Gamma_2$ .

**Proposition 4.1** Let  $\{\Gamma_j : j \in J\}$  be a family of NCTSs on  $X$ . Then  $\cap \Gamma_j$  is a neutrosophic crisp topology on  $X$ . Furthermore,  $\cap \Gamma_j$  is the coarsest NCT on  $X$  containing all topologies.

*Proof.* Obvious ■

Now, we define the neutrosophic crisp closure and neutrosophic crisp interior operations in neutrosophic crisp topological spaces:

**Definition 4.3** Let  $(X, \Gamma)$  be NCTS and  $A = \langle A_1, A_2, A_3 \rangle$  be a NCS in  $X$ . Then the neutrosophic crisp closure of  $A$  ( $NCCL(A)$  for short) and neutrosophic crisp interior ( $NCInt(A)$  for short) of  $A$  are defined by

$$\begin{aligned} NCCL(A) &= \cap \{K : \text{is an NCS in } X \text{ and } A \subseteq K\} \\ NCInt(A) &= \cup \{G : G \text{ is an NCOS in } X \text{ and } G \subseteq A\}, \end{aligned}$$

where NCS is a neutrosophic crisp set, and NCOS is a neutrosophic crisp open set.

It can be also shown that  $NCCL(A)$  is NCCS (neutrosophic crisp closed set) and  $NCInt(A)$  is a CNOS in  $X$ .

- a)  $A$  is in  $X$  if and only if  $NCCL(A) \supseteq A$ .
- b)  $A$  is an NCCS in  $X$  if and only if  $NCInt(A) = A$ .

**Proposition 4.2** For any neutrosophic crisp set  $A$  in  $(X, \Gamma)$  we have

- (a)  $NCCL(A^c) = (NCInt(A))^c$ .
- (b)  $NCInt(A^c) = (NCCL(A))^c$ .

*Proof.* Let  $A = \langle A_1, A_2, A_3 \rangle$  and suppose that the family of neutrosophic crisp subsets contained in  $A$  are indexed by the family if NCSs contained in  $A$  are indexed by the family  $A = \{ \langle A_{j1}, A_{j2}, A_{j3} \rangle : i \in J \}$ .

- a) Then we see that we have two types of

$$\begin{aligned} NCInt(A) &= \{ \langle \cup A_{j1}, \cup A_{j2}, \cap A_{j3} \rangle \} \text{ or} \\ NCInt(A) &= \{ \langle \cup A_{j1}, \cap A_{j2}, \cap A_{j3} \rangle \} \text{ hence} \\ (NCInt(A))^c &= \{ \langle \cap A_{j1}, \cap A_{j2}, \cup A_{j3} \rangle \} \text{ or} \\ (NCInt(A))^c &= \{ \langle \cap A_{j1}, \cup A_{j2}, \cup A_{j3} \rangle \}. \end{aligned}$$

- b) Hence  $NCCL(A^c) = (NCInt(A))^c$  follows immediately, which is analogous to (a). ■

**Proposition 4.3** Let  $(X, \Gamma)$  be a NCTS and  $A, B$  be two neutrosophic crisp sets in  $X$ . Then the following properties hold:

- (a)  $NCInt(A) \subseteq A$ ,
- (b)  $A \subseteq NCCL(A)$ ,
- (c)  $A \subseteq B \Rightarrow NCInt(A) \subseteq NCInt(B)$ ,
- (d)  $A \subseteq B \Rightarrow NCCL(A) \subseteq NCCL(B)$ ,
- (e)  $NCInt(A \cap B) = NCInt(A) \cap NCInt(B)$ ,
- (f)  $NCCL(A \cup B) = NCCL(A) \cup NCCL(B)$ ,
- (g)  $NCInt(X_N) = X_N$ ,
- (h)  $NCCL(\phi_N) = \phi_N$ .

*Proof.* (a), (b) and (e) are obvious; (c) follows from (a) and definitions. ■

## 5 Neutrosophic Crisp Continuity

Here come the basic definitions first:

**Definition 5.1** (a) If  $B = \langle B_1, B_2, B_3 \rangle$  is a NCS in  $Y$ , then the preimage of  $B$  under  $f$  denoted by  $f^{-1}(B)$  is a NCS in  $X$  defined by  $f^{-1}(B) = \langle f^{-1}(B_1), f^{-1}(B_2), f^{-1}(B_3) \rangle$ .

(b) If  $A = \langle A_1, A_2, A_3 \rangle$  is a NCS in  $X$ , then the image of  $A$  under  $f$  denoted by  $f(A)$  is the NCS in  $Y$  defined by  $f(A) = \langle f(A_1), f(A_2), f(A_3)^c \rangle$ .

Here we introduce the properties of images and preimages some of which we shall frequently use in the following sections.

**Corollary 5.1** Let  $A, \{A_i : i \in J\}$  be NCSs in  $X$ , and  $B, \{B_j : j \in K\}$  NCS in  $Y$ , and  $f : X \rightarrow Y$  a function. Then

- (a)  $A_1 \subseteq A_2 \Leftrightarrow f(A_1) \subseteq f(A_2)$ ,  
 $B_1 \subseteq B_2 \Leftrightarrow f^{-1}(B_1) \subseteq f^{-1}(B_2)$ ,
- (b)  $A \subseteq f^{-1}(f(A))$  and if  $f$  is injective, then  $A = f^{-1}(f(A))$ .
- (c)  $f^{-1}(f(B)) \subseteq B$  and if  $f$  is surjective, then  $f^{-1}(f(B)) = B$ .
- (d)  $f^{-1}(\cup B_i) = \cup f^{-1}(B_i)$ ,  $f^{-1}(\cap B_i) = \cap f^{-1}(B_i)$ ,
- (e)  $f(\cup A_i) = \cup f(A_i)$ ;  $f(\cap A_i) \subseteq \cap f(A_i)$ ; and if  $f$  is injective, then  $f(\cap A_i) = \cap f(A_i)$ ;
- (f)  $f^{-1}(Y_N) = X_N$ ,  $f^{-1}(\phi_N) = \phi_N$ .
- (g)  $f(\phi_N) = \phi_N$ ,  $f(X_N) = Y_N$ , if  $f$  is surjective.

*Proof.* Obvius. ■

**Definition 5.2** Let  $(X, \Gamma_1)$  and  $(Y, \Gamma_2)$  be two NCTSs, and let  $f : X \rightarrow Y$  be a function. Then  $f$  is said to be continuous iff the preimage of each NCS in  $\Gamma_2$  is an NCS in  $\Gamma_1$ .

**Definition 5.3** Let  $(X, \Gamma_1)$  and  $(Y, \Gamma_2)$  be two NCTSs and let  $f : X \rightarrow Y$  be a function. Then  $f$  is said to be open iff the image of each NCS in  $\Gamma_1$  is an NCS in  $\Gamma_2$ .

**Example 5.1** Let  $(X, \Gamma_0)$  and  $(Y, \psi_0)$  be two NCTSs.

(a) If  $f : X \rightarrow Y$  is continuous in the usual sense, then in this case,  $f$  is continuous in the sense of Definition 5.1 too. Here we consider the NCTs on  $X$  and  $Y$ , respectively, as follows:  $\Gamma_1 = \{\langle G, \phi, G^c \rangle : G \in \Gamma_0\}$  and  $\Gamma_2 = \{\langle H, \phi, H^c \rangle : H \in \Psi_0\}$ , In this case we have, for each  $\langle H, \phi, H^c \rangle \in \Gamma_2$ ,  $H \in \Psi_0$ ,  $f^{-1}\langle H, \phi, H^c \rangle = \langle f^{-1}(H), f^{-1}(\phi), f^{-1}(H^c) \rangle = \langle f^{-1}(H), f(\phi), (f(H))^c \rangle \in \Gamma_1$ .

(b) If  $f : X \rightarrow Y$  is open in the usual sense, then in this case,  $f$  is open in the sense of Definition 3.2.

Now we obtain some characterizations of continuity:

**Proposition 5.1** Let  $f : (X, \Gamma_1) \rightarrow (Y, \Gamma_2)$ .  $f$  is continuous iff the preimage of each CNCS (crisp neutrosophic closed set) in  $\Gamma_2$  is a CNCS in  $\Gamma_1$ .

**Proposition 5.2** The following are equivalent to each other:

- (a)  $f : (X, \Gamma_1) \rightarrow (Y, \Gamma_2)$  is continuous.
- (b)  $f^{-1}(CNInt(B)) \subseteq CNInt(f^{-1}(B))$  for each CNS  $B$  in  $Y$ .
- (c)  $CNCl(f^{-1}(B)) \subseteq f^{-1}(CNCl(B))$  for each CNC  $B$  in  $Y$ .

**Example 5.2** Let  $(Y, \Gamma_2)$  be an NCTS and  $f : X \rightarrow Y$  be a function. In this case  $\Gamma_1 = \{f^{-1}(H) : H \in \Gamma_2\}$  is a NCT on  $X$ . Indeed, it is the coarsest NCT on  $X$  which makes the function  $f : X \rightarrow Y$  continuous. One may call the initial neutrosophic crisp topology with respect to  $f$ .

## 6 Neutrosophic Crisp Compact Space (NCCS)

First we present the basic concepts:

**Definition 6.1** Let  $(X, \Gamma)$  be a NCTS.

- (a) If a family  $\{\langle G_{i_1}, G_{i_2}, G_{i_3} \rangle : i \in J\}$  of NCOSs in  $X$  satisfies the condition  $\cup\{\langle G_{i_1}, G_{i_2}, G_{i_3} \rangle : i \in J\} = X_N$  then it is called an neutrosophic open cover of  $X$ .
- (b) A finite subfamily of an open cover  $\{\langle G_{i_1}, G_{i_2}, G_{i_3} \rangle : i \in J\}$  on  $X$ , which is also a neutrosophic open cover of  $X$ , is called a neutrosophic finite subcover  $\{\langle G_{i_1}, G_{i_2}, G_{i_3} \rangle : i \in J\}$ .
- (c) A family  $\{\langle K_{i_1}, K_{i_2}, K_{i_3} \rangle : i \in J\}$  of NCCSs in  $X$  satisfies the finite intersection property (FIP for short) iff every finite subfamily  $\{\langle K_{i_1}, K_{i_2}, K_{i_3} \rangle : i = 1, 2, \dots, n\}$  of the family satisfies the condition  $\cap\{\langle K_{i_1}, K_{i_2}, K_{i_3} \rangle : i \in J\} \neq \phi_N$ .

**Definition 6.2** A NCTS  $(X, \Gamma)$  is called neutrosophic crisp compact iff each crisp neutrosophic open cover of  $X$  has a finite subcover.

**Example 6.1** (a) Let  $X = N$  and consider the NCSs given below:

$$\begin{aligned} A_1 &= \langle \{2, 3, 4, \dots\}, \phi, \phi \rangle, \\ A_2 &= \langle \{3, 4, 5, \dots\}, \phi, \{1\} \rangle, \\ A_3 &= \langle \{4, 5, 6, \dots\}, \phi, \{1, 2\} \rangle, \\ &\vdots \\ A_n &= \langle \{n+1, n+2, n+3, \dots\}, \phi, \{1, 2, 3, \dots, n-1\} \rangle. \end{aligned}$$

Then  $\Gamma = \{\phi_N, X_N\} \cup \{A_n = 3, 4, 5, \dots\}$  is an NCT on  $X$  and  $(X, \Gamma)$  is a neutrosophic crisp compact.

(b) Let  $X = (0, 1)$  and let's make the NCSs

$$A_n = \left\langle X, \left(\frac{1}{n}, \frac{n-1}{n}\right), \phi, \left(0, \frac{1}{n}\right) \right\rangle, \quad n = 3, 4, 5, \dots \text{ in } X$$

In this case  $\Gamma = \{\phi_N, X_n\} \cup \{A_n = 3, 4, 5, \dots\}$  is a NCT on  $X$ , which is not a neutrosophic crisp compact.

**Corollary 6.1** A NCTS  $(X, \Gamma)$  is neutrosophic crisp compact iff every family  $\{\langle X, G_{i_1}, G_{j_2}, G_{i_3} \rangle : i \in J\}$  of NCCSs in  $X$  having the FIP has nonempty intersection.

**Corollary 6.2** Let  $(X, \Gamma_1), (Y, \Gamma_2)$  be NCTSs and  $f : X \rightarrow Y$  be a continuous surjection. If  $(X, \Gamma_1)$  is a neutrosophic crisp compact, then so is  $(Y, \Gamma_2)$ .

**Definition 6.3** (a) If a family  $\{\langle X, G_{i_1}, G_{j_2}, G_{i_3} \rangle : i \in J\}$  of NCCSs in  $X$  satisfies the condition  $A \subseteq \cup\{\langle G_{i_1}, G_{j_2}, G_{i_3} \rangle : i \in J\}$ , then it is called a neutrosophic crisp open cover of  $A$ .

- (b) Let's consider a finite subfamily of a neutrosophic crisp open subcover of  $\{\langle X, G_{i_1}, G_{j_2}, G_{i_3} \rangle : i \in J\}$ . A neutrosophic crisp set  $A = \langle A_1, A_2, A_3 \rangle$  in a NCTS  $(X, \Gamma)$  is called neutrosophic crisp compact iff every neutrosophic crisp open cover of  $A$  has a finite neutrosophic crisp open subcover.

**Corollary 6.3** Let  $(X, \Gamma_1), (Y, \Gamma_2)$  be NCTSs and  $f : X \rightarrow Y$  is a continuous surjection. If  $A$  is a neutrosophic crisp compact in  $(X, \Gamma_1)$ , then so is  $f(A)$  in  $(Y, \Gamma_2)$ .

## 7 Conclusion

In this paper we introduced the neutrosophic crisp topology and the neutrosophic crisp compact space. Then we presented several properties for each of them.

## References

- [1] Florentin Smarandache, Neutrosophy and Neutro-sophic Logic, First International Conference on Neutrosophy, Neutrosophic Logic, Set, Probability, and Statistics, University of New Mexico, Gallup, NM 87301, USA(2002).
- [2] Florentin Smarandache, An introduction to the Neutrosophy probability applid in Quntum Physics, International Conference on introducation Neutro-soph Physics, Neutrosophic Logic, Set, Probabil-ity, and Statistics, University of New Mexico, Gallup, NM 87301, USA2-4 December (2011).
- [3] F. Smarandache, A Unifying Field in Logics: Neutro-sophic Logic. Neutrosophy, Neutrosophic Set, Neutro-sophic Probability. American Research Press, Reho-both, NM, (1999).
- [4] A.A. Salama and S.A. Alblowi, "Generalized Neutro-sophic Set and Generalized Neutrosophic Topologi-cal Spaces", Journal computer Sci. Engineering, Vol. (2) No. (7) (2012) pp 29–32.
- [5] A.A. Salama and S.A. Alblowi, Neutrosophic set and neutrosophic topological space, ISORJ. Mathemat-ics, Vol.(3), Issue(4), ( 2012), pp. 31–35.
- [6] A.A. Salama and S.A. Alblowi, Intuitionistic Fuzzy Ideals Topological Spaces, Advances in Fuzzy Math-ematics , Vol.(7), Number 1, (2012), pp. 51–60.
- [7] A.A. Salama, and H. Elagamy, "Neutrosophic Filters", International Journal of Computer Science Engineer-ing and Information Technology Reseach (IJCSEITR), Vol.3, Issue(1), Mar 2013, (2013), pp. 307–312.
- [8] S. A. Alblowi, A. A. Salama, and Mohmed Eisa, New Concepts of Neutrosophic Sets, International Journal of Mathematics and Computer Applications Research (IJMCAR), Vol.3, Issue 4, Oct 2013, (2013), pp. 95–102.
- [9] I. Hanafy, A.A. Salama and K. Mahfouz, Correlation of neutrosophic Data, International Refereed Journal of Engineering and Science (IRJES) , Vol.(1), Issue 2 , (2012), pp. 39–43.
- [10] I.M. Hanafy, A.A. Salama and K.M. Mahfouz, "Neutrosophic Crisp Events and Its Probability" Inter-national Journal of Mathematics and Computer Ap-plications Research (IJMCAR) Vol.(3), Issue 1, Mar 2013, (2013), pp. 171–178.
- [11] A. A. Salama, "Neutrosophic Crisp Points & Neutrosophic Crisp Ideals", Neutrosophic Sets and Systems, Vol.1, No. 1, (2013) pp. 50–54.
- [12] A. A. Salama and F. Smarandache, "Filters via Neutrosophic Crisp Sets", Neutrosophic Sets and Systems, Vol.1, No. 1, (2013) pp. 34–38.