

Some Types of Neutrosophic Crisp Sets and Neutrosophic Crisp Relations

A. A. Salama¹, Said Broumi² and Florentin Smarandache³

Department of Mathematics and Computer Science, Faculty of Science, Port Said University, Egypt

drsalama44@gmail.com

²Faculty of Arts and Humanities, Hay El Baraka Ben M'sik Casablanca B.P. 7951, Hassan II Casablanca University, Morocco.

broumisaid78@gmail.com

³University of New Mexico, 705 Gurley Ave., Gallup, New Mexico 87301, USA

fsmarandache@gmail.com

Abstract—The purpose of this paper is to introduce a new types of crisp sets are called the neutrosophic crisp set with three types 1, 2, 3. After given the fundamental definitions and operations, we obtain several properties, and discussed the relationship between neutrosophic crisp sets and others. Finally, we introduce and study the notion of neutrosophic crisp relations.

Index Terms—Neutrosophic set, neutrosophic crisp sets, neutrosophic crisp relations, generalized neutrosophic set, Intuitionistic neutrosophic Set.

I. Introduction

Since the world is full of indeterminacy, the neutrosophics found their place into contemporary research. The fundamental concepts of neutrosophic set, introduced by Smarandache in [16, 17, 18], and Salama et al. in [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15], provides a natural foundation for treating mathematically the neutrosophic phenomena which exist pervasively in our real world and for building new branches of neutrosophic mathematics. Neutrosophy has laid the foundation for a whole family of new mathematical theories generalizing both their classical and fuzzy counterparts [1, 2, 3, 19] such as a neutrosophic set theory. In this paper we introduce a new types of crisp sets are called the neutrosophic crisp set with three types 1, 2, 3. After given the fundamental definitions and operations, we obtain several properties, and discussed the relationship between

neutrosophic crisp sets and others. Finally, we introduce and study the notion of neutrosophic crisp relations.

The paper unfolds as follows. The next section briefly introduces some definitions related to neutrosophic set theory and some terminologies of neutrosophic crisp set. Section 3 presents new types of neutrosophic crisp sets and studied some of their basic properties. Section 4 presents the concept of neutrosophic crisp relations. Finally we concludes the paper.

II. Preliminaries

We recollect some relevant basic preliminaries, and in particular, the work of Smarandache in [16, 17, 18], and Salama et al. [4,5]. Smarandache introduced the neutrosophic components T, I, F which represent the membership, indeterminacy, and non-membership values respectively, where $]^{-}0, 1^{+}[$ is nonstandard unit interval.

Definition 2.1 [9, 13, 15]

A neutrosophic crisp set (NCS for short)

$A = \langle A_1, A_2, A_3 \rangle$ can be identified to an ordered triple $\langle A_1, A_2, A_3 \rangle$ are subsets on X , and every crisp event in X is obviously an NCS having the form $\langle A_1, A_2, A_3 \rangle$,

Salama et al. constructed the tools for developed neutrosophic crisp set, and introduced the NCS ϕ_N, X_N in X as follows:

1) ϕ_N may be defined as four types:

i) Type1: $\phi_N = \langle \phi, \phi, X \rangle$, or

ii) Type2: $\phi_N = \langle \phi, X, X \rangle$, or

iii) Type3: $\phi_N = \langle \phi, X, \phi \rangle$, or

iv) Type4: $\phi_N = \langle \phi, \phi, \phi \rangle$

2) X_N may be defined as four types

i) Type1: $X_N = \langle X, \phi, \phi \rangle$,

ii) Type2: $X_N = \langle X, X, \phi \rangle$,

iii) Type3: $X_N = \langle X, X, X \rangle$,

iv) Type4: $X_N = \langle X, X, X \rangle$,

Definition 2.2 [9, 13, 15]

Let $A = \langle A_1, A_2, A_3 \rangle$ a NCE or UNCE on X , then

the complement of the set A (A^c , for short) maybe defined as three kinds of complements

(C₁) Type1: $A^c = \langle A^c_1, A^c_2, A^c_3 \rangle$,

(C₂) Type2: $A^c = \langle A_3, A_2, A_1 \rangle$

(C₃) Type3: $A^c = \langle A_3, A^c_2, A_1 \rangle$

One can define several relations and operations between NCS as follows:

Definition 2.3 [9, 13, 15]

Let X be a non-empty set, and NCSS A and B in the form $A = \langle A_1, A_2, A_3 \rangle, B = \langle B_1, B_2, B_3 \rangle$, then we may consider two possible definitions for subsets ($A \subseteq B$)

($A \subseteq B$) may be defined as two types:

1) Type1:

$$A \subseteq B \Leftrightarrow A_1 \subseteq B_1, A_2 \subseteq B_2 \text{ and } A_3 \supseteq B_3 \text{ or}$$

2) Type2:

$$A \subseteq B \Leftrightarrow A_1 \subseteq B_1, A_2 \supseteq B_2 \text{ and } A_3 \supseteq B_3$$

Definition 2.4 [9, 13, 15]

Let X be a non-empty set, and NCSS A and B in the form $A = \langle A_1, A_2, A_3 \rangle, B = \langle B_1, B_2, B_3 \rangle$ are NCSS

Then

1) $A \cap B$ may be defined as two types:

i) Type1:

$$A \cap B = \langle A_1 \cap B_1, A_2 \cap B_2, A_3 \cup B_3 \rangle \text{ or}$$

ii) Type2:

$$A \cap B = \langle A_1 \cap B_1, A_2 \cup B_2, A_3 \cup B_3 \rangle$$

2) $A \cup B$ may be defined as two types:

i) Type 1: $A \cup B = \langle A_1 \cup B_1, A_2 \cap B_2, A_3 \cup B_3 \rangle$

or

ii) Type 2: $A \cup B = \langle A_1 \cup B_1, A_2 \cap B_2, A_3 \cap B_3 \rangle$

Proposition 2.1 [9, 13, 15]

Let $\{A_j : j \in J\}$ be arbitrary family of neutrosophic crisp subsets in X , then

1) $\cap A_j$ may be defined two types as :

i) Type1: $\cap A_j = \langle \cap A_{j_1}, \cap A_{j_2}, \cup A_{j_3} \rangle$, or

ii) Type2: $\cap A_j = \langle \cap A_{j_1}, \cup A_{j_2}, \cup A_{j_3} \rangle$.

2) $\cup A_j$ may be defined two types as :

i) Type1: $\cup A_j = \langle \cup A_{j_1}, \cap A_{j_2}, \cap A_{j_3} \rangle$ or

ii) Type2: $\cup A_j = \langle \cup A_{j_1}, \cup A_{j_2}, \cap A_{j_3} \rangle$.

III. New Types of Neutrosophic Crisp Sets

We shall now consider some possible definitions for some types of neutrosophic crisp sets

Definition 3.1

Let X be a non-empty fixed sample space. A neutrosophic crisp set (NCS for short) A is an object having the form $A = \langle A_1, A_2, A_3 \rangle$ where A_1, A_2 and A_3 are subsets of X .

Definition 3.2

The object having the form $A = \langle A_1, A_2, A_3 \rangle$ is called

1) (Neutrosophic Crisp Set with Type 1) If

$$\text{satisfying } A_1 \cap A_2 = \phi, A_1 \cap A_3 = \phi$$

$$\text{and } A_2 \cap A_3 = \phi. \text{ (NCS-Type1 for short).}$$

2) (Neutrosophic Crisp Set with Type 2) If

$$\text{satisfying } A_1 \cap A_2 = \phi, A_1 \cap A_3 = \phi$$

$$\text{and } A_2 \cap A_3 = \phi \text{ and } A_1 \cup A_2 \cup A_3 = X. \text{ (NCS-Type2 for short).}$$

3) (Neutrosophic Crisp Set with Type 3) If satisfying
 $A_1 \cap A_2 \cap A_3 = \emptyset$
 $A_1 \cup A_2 \cup A_3 = X$. (NCS-Type3 for short).

Definition 3.3

1) (Neutrosophic Set [7]): Let X be a non-empty fixed set. A neutrosophic set (NS for short) A is an object having the form $A = \langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle$ where $\mu_A(x)$, $\sigma_A(x)$ and $\nu_A(x)$ which represent the degree of member ship function (namely $\mu_A(x)$), the degree of indeterminacy (namely $\sigma_A(x)$), and the degree of non-member ship (namely $\nu_A(x)$) respectively of each element $x \in X$ to the set A where $0^- \leq \mu_A(x), \sigma_A(x), \nu_A(x) \leq 1^+$ and $0^- \leq \mu_A(x) + \sigma_A(x) + \nu_A(x) \leq 3^+$.

2) (Generalized Neutrosophic Set [8]): Let X be a non-empty fixed set. A generalized neutrosophic (GNS for short) set A is an object having the form $A = \langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle$ where $\mu_A(x)$, $\sigma_A(x)$ and $\nu_A(x)$ which represent the degree of member ship function (namely $\mu_A(x)$), the degree of indeterminacy (namely $\sigma_A(x)$), and the degree of non-member ship (namely $\nu_A(x)$) respectively of each element $x \in X$ to the set A where $0^- \leq \mu_A(x), \sigma_A(x), \nu_A(x) \leq 1^+$ and the functions satisfy the condition $\mu_A(x) \wedge \sigma_A(x) \wedge \nu_A(x) \leq 0.5$ and $0^- \leq \mu_A(x) + \sigma_A(x) + \nu_A(x) \leq 3^+$.

3) (Intuitionistic Neutrosophic Set [16]). Let X be a non-empty fixed set. An intuitionistic neutrosophic set A (INS for short) is an object having the form $A = \langle \mu_A(x), \sigma_A(x), \nu_A(x) \rangle$ where $\mu_A(x)$, $\sigma_A(x)$ and $\nu_A(x)$ which represent the degree of member ship function (namely $\mu_A(x)$), the degree of indeterminacy (namely $\sigma_A(x)$), and the degree of non-member ship (namely $\nu_A(x)$) respectively of each element $x \in X$ to the set A where $0.5 \leq \mu_A(x), \sigma_A(x), \nu_A(x)$ and the functions satisfy the condition $\mu_A(x) \wedge \sigma_A(x) \leq 0.5$,

$$\mu_A(x) \wedge \nu_A(x) \leq 0.5, \quad \sigma_A(x) \wedge \nu_A(x) \leq 0.5, \text{ and } 0 \leq \mu_A(x) + \sigma_A(x) + \nu_A(x) \leq 2^+.$$

A neutrosophic crisp with three types the object $A = \langle A_1, A_2, A_3 \rangle$ can be identified to an ordered triple $\langle A_1, A_2, A_3 \rangle$ are subsets on X, and every crisp set in X is obviously a NCS having the form $\langle A_1, A_2, A_3 \rangle$. Every neutrosophic set $A = \langle \mu_A(x), \sigma_A(x), \nu_A(x) \rangle$ on X is obviously on NS having the form $\langle \mu_A(x), \sigma_A(x), \nu_A(x) \rangle$.

Remark 3.1

- 1) The neutrosophic set not to be generalized neutrosophic set in general.
- 2) The generalized neutrosophic set in general not intuitionistic NS but the intuitionistic NS is generalized NS.

Intuitionistic NS \longrightarrow Generalized NS \longrightarrow NS




Fig.1: Represents the relation between types of NS

Corollary 3.1

Let X non-empty fixed set and $A = \langle \mu_A(x), \sigma_A(x), \nu_A(x) \rangle$ be INS on X
 Then:

- 1) Type1- A^c of INS be a GNS.
- 2) Type2- A^c of INS be a INS.
- 3) Type3- A^c of INS be a GNS.

Proof

Since A INS then $0.5 \leq \mu_A(x), \sigma_A(x), \nu_A(x)$, and $\mu_A(x) \wedge \sigma_A(x) \leq 0.5, \nu_A(x) \wedge \mu_A(x) \leq 0.5, \nu_A(x) \wedge \sigma_A(x) \leq 0.5$ Implies $\mu^c_A(x), \sigma^c_A(x), \nu^c_A(x) \leq 0.5$ then is not to be Type1- A^c INS. On other hand the Type 2- A^c , $A^c = \langle \nu_A(x), \sigma_A(x), \mu_A(x) \rangle$ be INS and Type 3- A^c , $A^c = \langle \nu_A(x), \sigma^c_A(x), \mu_A(x) \rangle$ and $\sigma^c_A(x) \leq 0.5$ implies to $A^c = \langle \nu_A(x), \sigma^c_A(x), \mu_A(x) \rangle$ GNS and not to be INS

Example 3.1

Let $X = \{a, b, c\}$, and A, B, C are neutrosophic sets on X ,

$$A = \langle 0.7, 0.9, 0.8 \rangle \setminus a, \langle 0.6, 0.7, 0.6 \rangle \setminus b, \langle 0.9, 0.7, 0.8 \rangle \setminus c,$$

$$B = \langle 0.7, 0.9, 0.5 \rangle \setminus a, \langle 0.6, 0.4, 0.5 \rangle \setminus b, \langle 0.9, 0.5, 0.8 \rangle \setminus c$$

$$C = \langle 0.7, 0.9, 0.5 \rangle \setminus a, \langle 0.6, 0.8, 0.5 \rangle \setminus b, \langle 0.9, 0.5, 0.8 \rangle \setminus c$$

By the Definition 3.3 no.3

$\mu_A(x) \wedge \sigma_A(x) \wedge \nu_A(x) \geq 0.5$, A be not GNS and INS,

$$B = \langle 0.7, 0.9, 0.5 \rangle \setminus a, \langle 0.6, 0.4, 0.5 \rangle \setminus b, \langle 0.9, 0.5, 0.8 \rangle \setminus c$$

not INS, where $\sigma_A(b) = 0.4 < 0.5$. Since

$\mu_B(x) \wedge \sigma_B(x) \wedge \nu_B(x) \leq 0.5$ then B is a GNS but not INS.

$$A^c = \langle 0.3, 0.1, 0.2 \rangle \setminus a, \langle 0.4, 0.3, 0.4 \rangle \setminus b, \langle 0.1, 0.3, 0.2 \rangle \setminus c$$

be a GNS, but not INS.

$$B^c = \langle 0.3, 0.1, 0.5 \rangle \setminus a, \langle 0.4, 0.6, 0.5 \rangle \setminus b, \langle 0.1, 0.5, 0.2 \rangle \setminus c$$

be a GNS, but not INS, C be INS and GNS,

$$C^c = \langle 0.3, 0.1, 0.5 \rangle \setminus a, \langle 0.4, 0.2, 0.5 \rangle \setminus b, \langle 0.1, 0.5, 0.2 \rangle \setminus c$$

be a GNS but not INS.

Definition 3.4

A neutrosophic crisp set (NCS for short)

$A = \langle A_1, A_2, A_3 \rangle$ can be identified to an ordered triple $\langle A_1, A_2, A_3 \rangle$ are subsets on X , and every crisp set in X is obviously an NCS having the form $\langle A_1, A_2, A_3 \rangle$,

Salama et al in [6,13] constructed the tools for developed neutrosophic crisp set, and introduced the NCS ϕ_N, X_N in X as follows:

1) ϕ_N may be defined as four types:

i) Type1: $\phi_N = \langle \phi, \phi, X \rangle$, or

ii) Type2: $\phi_N = \langle \phi, X, X \rangle$, or

iii) Type3: $\phi_N = \langle \phi, X, \phi \rangle$, or

iv) Type4: $\phi_N = \langle \phi, \phi, \phi \rangle$

2) X_N may be defined as four types

i) Type1: $X_N = \langle X, \phi, \phi \rangle$,

ii) Type2: $X_N = \langle X, X, \phi \rangle$,

v) Type3: $X_N = \langle X, X, X \rangle$,

vi) Type4: $X_N = \langle X, X, X \rangle$,

Definition 3.5

A NCS-Type1 ϕ_{N1}, X_{N1} in X as follows:

1) ϕ_{N1} may be defined as three types:

i) Type1: $\phi_{N1} = \langle \phi, \phi, X \rangle$, or

ii) Type2: $\phi_{N1} = \langle \phi, X, \phi \rangle$, or

iii) Type3: $\phi_N = \langle \phi, \phi, \phi \rangle$.

2) X_{N1} may be defined as one type

Type1: $X_{N1} = \langle X, \phi, \phi \rangle$.

Definition 3.6

A NCS-Type2, ϕ_{N2}, X_{N2} in X as follows:

1) ϕ_{N2} may be defined as two types:

i) Type1: $\phi_{N2} = \langle \phi, \phi, X \rangle$, or

ii) Type2: $\phi_{N2} = \langle \phi, X, \phi \rangle$

2) X_{N2} may be defined as one type

Type1: $X_{N2} = \langle X, \phi, \phi \rangle$

Definition 3.7

a NCS-Type 3, ϕ_{N3}, X_{N3} in X as follows:

1) ϕ_{N3} may be defined as three types:

i) Type1: $\phi_{N3} = \langle \phi, \phi, X \rangle$, or

ii) Type2: $\phi_{N3} = \langle \phi, X, \phi \rangle$, or

iii) Type3: $\phi_{N3} = \langle \phi, X, X \rangle$.

2) X_{N3} may be defined as three types

i) Type1: $X_{N3} = \langle X, \phi, \phi \rangle$,

ii) Type2: $X_{N3} = \langle X, X, \phi \rangle$,

iii) Type3: $X_{N3} = \langle X, \phi, X \rangle$,

Corollary 3.1

In general

1-Every NCS-Type 1, 2, 3 are NCS.

2-Every NCS-Type 1 not to be NCS-Type2, 3.

3-Every NCS-Type 2 not to be NCS-Type1, 3.

4-Every NCS-Type 3 not to be NCS-Type2, 1, 2.

5-Every crisp set be NCS.

The following Venn diagram represents the relation between NCSs




Fig. 2: Venn diagram represents the relation between NCSs

Example 3.2

Let A, B, C, D are NCSs on $X = \{a, b, c, d, e, f\}$, the following types of neutrosophic crisp sets

- i) $A = \langle \{a\}, \{b\}, \{c\} \rangle$ be a NCS-Type 1, but not NCS-Type 2 and Type 3
- ii) $B = \langle \{a, b\}, \{c, d\}, \{f, e\} \rangle$ be a NCS-Type 1, 2, 3
- iii) $C = \langle \{a, b, c, d\}, \{e\}, \{a, b, f\} \rangle$ be a NCS-Type 3 but not NCS-Type 1, 2.
- iv) $D = \langle \{a, b, c, d\}, \{a, b, c\}, \{a, b, d, f\} \rangle$ be a NCS but not NCS-Type 1, 2, 3.

The complement for A, B, C, D may be equals
The complement of A

- i)Type 1: $A^c = \langle \{b, c, d, e, f\}, \{a, c, d, e, f\}, \{a, b, d, e, f\} \rangle$ be a NCS but not NCS—Type1, 2,3
- ii)Type 2: $A^c = \langle \{c\}, \{b\}, \{a\} \rangle$ be a NCS-Type 3 but not NCS—Type1, 2
- iii)Type 3: $A^c = \langle \{c\}, \{a, c, d, e, f\}, \{a\} \rangle$ be a NCS-Type 1 but not NCS—Type 2, 3.

The complement of B may be equals

- i)Type 1: $B^c = \langle \{c, d, e, f\}, \{a, b, e, f\}, \{a, b, c, d\} \rangle$ be NCS-Type 3 but not NCS-Type 1, 2.
- ii) Type 2: $B^c = \langle \{e, f\}, \{c, d\}, \{a, b\} \rangle$ be NCS-Type 1, 2, 3.
- iii)Type 3: $B^c = \langle \{e, f\}, \{a, b, e, f\}, \{a, b\} \rangle$ be NCS-Type 3, but not NCS-Type 1, 2.

The complement of C may be equals

- i)Type 1: $C^c = \langle \{e, f\}, \{a, b, c, d, f\}, \{c, d, e\} \rangle$.
- ii)Type 2: $C^c = \langle \{a, b, f\}, \{e\}, \{a, b, c, d\} \rangle$,
- iii)Type 3: $C^c = \langle \{a, b, f\}, \{a, b, c, d\}, \{a, b, c, d\} \rangle$,

The complement of D may be equals

- i)Type 1: $D^c = \langle \{e, f\}, \{d, e, f\}, \{c, e\} \rangle$ be NCS-Type 3 but not NCS-Type 1, 2.
- ii)Type 2: $D^c = \langle \{a, b, d, f\}, \{a, b, c\}, \{a, b, c, d\} \rangle$ be NCS-Type 3 but not NCS-Type 1, 2.
- iii)Type 3: $D^c = \langle \{a, b, d, f\}, \{d, e, f\}, \{a, b, c, d\} \rangle$ be NCS-Type 3 but not NCS-Type 1, 2.

Definition 3.8

Let X be a non-empty set, $A = \langle A_1, A_2, A_3 \rangle$

1) If A be a NCS-Type1 on X , then the complement of the set A (A^c , for short) maybe defined as one kind of complement Type1:

$$A^c = \langle A_3, A_2, A_1 \rangle .$$

2) If A be a NCS-Type 2 on X , then the complement of the set A (A^c , for short) maybe defined as one kind of complement $A^c = \langle A_3, A_2, A_1 \rangle$.

3)If A be NCS-Type3 on X , then the complement of the set A (A^c , for short) maybe defined as one kind of complement defined as three kinds of complements

- (C_1) Type1: $A^c = \langle A^c_1, A^c_2, A^c_3 \rangle$,
- (C_2) Type2: $A^c = \langle A_3, A_2, A_1 \rangle$
- (C_3) Type3: $A^c = \langle A_3, A^c_2, A_1 \rangle$

Example 3.3

Let $X = \{a, b, c, d, e, f\}$, $A = \langle \{a, b, c, d\}, \{e\}, \{f\} \rangle$ be a NCS-Type 2, $B = \langle \{a, b, c\}, \{\emptyset\}, \{d, e\} \rangle$ be a NCS-Type1., $C = \langle \{a, b\}, \{c, d\}, \{e, f\} \rangle$ NCS-Type 3, then the complement $A = \langle \{a, b, c, d\}, \{e\}, \{f\} \rangle$, $A^c = \langle \{f\}, \{e\}, \{a, b, c, d\} \rangle$ NCS-Type 2, the complement of $B = \langle \{a, b, c\}, \{\emptyset\}, \{d, e\} \rangle$, $B^c = \langle \{d, e\}, \{\emptyset\}, \{a, b, c\} \rangle$ NCS-Type1. The complement of $C = \langle \{a, b\}, \{c, d\}, \{e, f\} \rangle$ may be defined as three types:

- Type 1: $C^c = \langle \{c, d, e, f\}, \{a, b, e, f\}, \{a, b, c, d\} \rangle$.
- Type 2: $C^c = \langle \{e, f\}, \{c, d\}, \{a, b\} \rangle$,
- Type 3: $C^c = \langle \{e, f\}, \{a, b, e, f\}, \{a, b\} \rangle$,

Proposition 3.1

Let $\{A_j : j \in J\}$ be arbitrary family of neutrosophic crisp subsets on X , then

- 1) $\cap A_j$ may be defined two types as :

Type1: $\cap A_j = \langle \cap A_{j_1}, \cap A_{j_2}, \cup A_{j_3} \rangle$, or

Type2: $\cap A_j = \langle \cap A_{j_1}, \cup A_{j_2}, \cup A_{j_3} \rangle$.

2) $\cup A_j$ may be defined two types as :

Type1: $\cup A_j = \langle \cup A_{j_1}, \cap A_{j_2}, \cap A_{j_3} \rangle$ or

Type2: $\cup A_j = \langle \cup A_{j_1}, \cup A_{j_2}, \cap A_{j_3} \rangle$.

Definition 3.9

(a) If $B = \langle B_1, B_2, B_3 \rangle$ is a NCS in Y, then the preimage of B under f , denoted by $f^{-1}(B)$, is a NCS in X defined by $f^{-1}(B) = \langle f^{-1}(B_1), f^{-1}(B_2), f^{-1}(B_3) \rangle$.

(b) If $A = \langle A_1, A_2, A_3 \rangle$ is a NCS in X, then the image of A under f , denoted by $f(A)$, is the a NCS in Y defined by $f(A) = \langle f(A_1), f(A_2), f(A_3)^c \rangle$.

Here we introduce the properties of images and preimages some of which we shall frequently use in the following.

Corollary 3.2

Let A, $\{A_i : i \in J\}$, be a family of NCS in X, and B, $\{B_j : j \in K\}$ NCS in Y, and $f : X \rightarrow Y$ a function. Then

- (a) $A_1 \subseteq A_2 \Leftrightarrow f(A_1) \subseteq f(A_2)$,
- $B_1 \subseteq B_2 \Leftrightarrow f^{-1}(B_1) \subseteq f^{-1}(B_2)$,
- (b) $A \subseteq f^{-1}(f(A))$ and if f is injective, then $A = f^{-1}(f(A))$,
- (c) $f^{-1}(f(B)) \subseteq B$ and if f is surjective, then $f^{-1}(f(B)) = B$,
- (d) $f^{-1}(\cup B_i) = \cup f^{-1}(B_i)$, $f^{-1}(\cap B_i) = \cap f^{-1}(B_i)$,
- (e) $f(\cup A_i) = \cup f(A_i)$; $f(\cap A_i) \subseteq \cap f(A_i)$; and if f is injective, then $f(\cap A_i) = \cap f(A_i)$;
- (f) $f^{-1}(Y_N) = X_N$, $f^{-1}(\phi_N) = \phi_N$.
- (g) $f(\phi_N) = \phi_N$, $f(X_N) = Y_N$, if f is subjective.

Proof

Obvious

IV. Neutrosophic Crisp Relations

Here we give the definition relation on neutrosophic crisp sets and study of its properties.

Let X, Y and Z be three ordinary nonempty sets

Definition 4.1

Let X and Y are two non-empty crisp sets and NCSS A and B in the form $A = \langle A_1, A_2, A_3 \rangle$ on X,

$B = \langle B_1, B_2, B_3 \rangle$ on Y. Then

i) The product of two neutrosophic crisp sets A and B is a neutrosophic crisp set $A \times B$ given by

$A \times B = \langle A_1 \times B_1, A_2 \times B_2, A_3 \times B_3 \rangle$ on $X \times Y$.

ii) We will call a neutrosophic crisp relation $R \subseteq A \times B$ on the direct product $X \times Y$.

The collection of all neutrosophic crisp relations on $X \times Y$ is denoted as $NCR(X \times Y)$

Definition 4.2

Let R be a neutrosophic crisp relation on $X \times Y$, then the inverse of R is denoted by R^{-1} where $R \subseteq A \times B$ on $X \times Y$ then $R^{-1} \subseteq B \times A$ on $Y \times X$.

Example 4.1

Let $X = \{a, b, c, d\}$, $A = \langle \{a, b\}, \{c\}, \{d\} \rangle$ and

$B = \langle \{a\}, \{c\}, \{d, b\} \rangle$ then the product of two neutrosophic crisp sets given by

$A \times B = \langle \{(a, a), (b, a)\}, \{(c, c)\}, \{(d, d), (d, b)\} \rangle$

and

$B \times A = \langle \{(a, a), (a, b)\}, \{(c, c)\}, \{(d, d), (b, d)\} \rangle$,

and

$R_1 = \langle \{(a, a)\}, \{(c, c)\}, \{(d, d)\} \rangle$, $R_1 \subseteq A \times B$ on $X \times X$,

$R_2 = \langle \{(a, b)\}, \{(c, c)\}, \{(d, d), (b, d)\} \rangle$

$R_2 \subseteq B \times A$ on $X \times X$.

Example 4.2

From the Example 3.1

$R_1^{-1} = \langle \{(a, a)\}, \{(c, c)\}, \{(d, d)\} \rangle \subseteq B \times A$ and

$R_2^{-1} = \langle \{(b, a)\}, \{(c, c)\}, \{(d, d), (d, b)\} \rangle$

$\subseteq B \times A$.

Example 4.3

Let $X = \{a, b, c, d, e, f\}$,

$A = \langle \{a, b, c, d\}, \{e\}, \{f\} \rangle$,

$D = \langle \{a, b\}, \{e, c\}, \{f, d\} \rangle$ be a NCS-Type 2,

$B = \langle \{a, b, c\}, \{\phi\}, \{d, e\} \rangle$ be a NCS-Type1.

$C = \langle \{a, b\}, \{c, d\}, \{e, f\} \rangle$ be a NCS-Type 3. Then

$$A \times D = \left\langle \begin{array}{l} \{(a, a), (a, b), (b, a), (b, b), (b, b), (c, a), (c, b)\} \\ \{(d, a), (d, b)\}, \{(e, e), (e, c)\}, \{(f, f), (f, d)\} \end{array} \right\rangle$$

$$D \times C = \left\langle \begin{array}{l} \{(a, a), (a, b), (b, a), (b, b)\}, \{(c, c), (c, d), \\ (c, c), (c, d)\}, \{(f, e), (f, f), (d, e), (d, f)\} \end{array} \right\rangle$$

we can construct many types of relations on products. We can define the operations of neutrosophic crisp relation.

Definition 4.4

Let R and S be two neutrosophic crisp relations between X and Y for every $(x, y) \in X \times Y$ and NCSS A and B in the form $A = \langle A_1, A_2, A_3 \rangle$ on X , $B = \langle B_1, B_2, B_3 \rangle$ on Y , Then we can defined the following operations

i) $R \subseteq S$ may be defined as two types

a) Type1: $R \subseteq S \Leftrightarrow A_{1R} \subseteq B_{1S}, A_{2R} \subseteq B_{2S},$

$$A_{3R} \supseteq B_{3S}$$

b) Type2: $R \subseteq S \Leftrightarrow A_{1R} \subseteq B_{1S}, A_{2R} \supseteq B_{2S},$

$$B_{3S} \subseteq A_{3R}$$

ii) $R \cup S$ may be defined as two types

a) Type1:

$$R \cup S = \langle A_{1R} \cup B_{1S}, A_{2R} \cup B_{2S}, A_{3R} \cap B_{3S} \rangle,$$

b) Type2:

$$R \cup S = \langle A_{1R} \cup B_{1S}, A_{2R} \cap B_{2S}, A_{3R} \cap B_{3S} \rangle.$$

iii) $R \cap S$ may be defined as two types

a) Type1: $R \cap S$

$$= \langle A_{1R} \cap B_{1S}, A_{2R} \cup B_{2S}, A_{3R} \cup B_{3S} \rangle,$$

b) Type2:

$$R \cap S = \langle A_{1R} \cap B_{1S}, A_{2R} \cap B_{2S}, A_{3R} \cup B_{3S} \rangle.$$

Theorem 4.1

Let R, S and Q be three neutrosophic crisp relations between X and Y for every $(x, y) \in X \times Y$, then

i) $R \subseteq S \Rightarrow R^{-1} \subseteq S^{-1}.$

ii) $(R \cup S)^{-1} \Rightarrow R^{-1} \cup S^{-1}.$

iii) $(R \cap S)^{-1} \Rightarrow R^{-1} \cap S^{-1}.$

iv) $(R^{-1})^{-1} = R.$

v) $R \cap (S \cup Q) = (R \cap S) \cup (R \cap Q).$

vi) $R \cup (S \cap Q) = (R \cup S) \cap (R \cup Q).$

vii) If $S \subseteq R, Q \subseteq R$, then $S \cup Q \subseteq R$

Proof

Clear

Definition 4.5

The neutrosophic crisp relation $I \in NCR(X \times X)$, the neutrosophic crisp relation of identity may be defined as two types

i) Type1: $I = \langle \{A \times A\}, \{A \times A\}, \phi \rangle$

ii) Type2: $I = \langle \{A \times A\}, \phi, \phi \rangle$

Now we define two composite relations of neutrosophic crisp sets.

Definition 4.6

Let R be a neutrosophic crisp relation in $X \times Y$, and S be a neutrosophic crisp relation in $Y \times Z$. Then the composition of R and S , $R \circ S$ be a neutrosophic crisp relation in $X \times Z$ as a definition may be defined as two types

i) Type1:

$$R \circ S \leftrightarrow (R \circ S)(x, z)$$

$$= \cup \{ \langle \{ (A_1 \times B_1)_R \cap (A_2 \times B_2)_S \}, \{ (A_2 \times B_2)_R \cap (A_2 \times B_2)_S \}, \{ (A_3 \times B_3)_R \cap (A_3 \times B_3)_S \} \rangle \}.$$

ii) Type2 :

$$R \circ S \leftrightarrow (R \circ S)(x, z)$$

$$= \cap \{ \langle \{ (A_1 \times B_1)_R \cup (A_2 \times B_2)_S \}, \{ (A_2 \times B_2)_R \cup (A_2 \times B_2)_S \}, \{ (A_3 \times B_3)_R \cup (A_3 \times B_3)_S \} \rangle \}.$$

Example 4.5

Let $X = \{a, b, c, d\}$, $A = \langle \{a, b\}, \{c\}, \{d\} \rangle$ and

$B = \langle \{a\}, \{c\}, \{d, b\} \rangle$ then the product of two events given

by $A \times B = \langle \{ (a, a), (b, a) \}, \{ (c, c) \}, \{ (d, d), (d, b) \} \rangle,$

and

$B \times A = \langle \{ (a, a), (a, b) \}, \{ (c, c) \}, \{ (d, d), (b, d) \} \rangle,$

and

$R_1 = \langle \{ (a, a) \}, \{ (c, c) \}, \{ (d, d) \} \rangle, R_1 \subseteq A \times B$ on $X \times X$,

$R_2 = \langle \{ (a, b) \}, \{ (c, c) \}, \{ (d, d), (b, d) \} \rangle$

$R_2 \subseteq B \times A$ on $X \times X$.

$$R_1 \circ R_2 = \cup \langle \{(a, a)\} \cap \{(a, b)\}, \{(c, c)\}, \{(d, d)\} \rangle$$

$$= \langle \{\emptyset\}, \{(c, c)\}, \{(d, d)\} \rangle \text{ and}$$

$$I_{A1} = \langle \{(a, a).(a, b).(b, a)\}, \{(a, a).(a, b).(b, a)\}, \{\emptyset\} \rangle$$

$$, I_{A2} = \langle \{(a, a).(a, b).(b, a)\}, \{\emptyset\}, \{\emptyset\} \rangle$$

Theorem 4.2

Let R be a neutrosophic crisp relation in $X \times Y$, and S be a neutrosophic crisp relation in $Y \times Z$ then $(R \circ S)^{-1} = S^{-1} \circ R^{-1}$.

Proof

Let $R \subseteq A \times B$ on $X \times Y$ then $R^{-1} \subseteq B \times A$,
 $S \subseteq B \times D$ on $Y \times Z$ then $S^{-1} \subseteq D \times B$, from
 Definition 3.6 and similarly we
 can $I_{(R \circ S)^{-1}}(x, z) = I_{S^{-1}}(x, z)$ and $I_{R^{-1}}(x, z)$ then
 $(R \circ S)^{-1} = S^{-1} \circ R^{-1}$.

V. Conclusion

In our work, we have put forward some new types of neutrosophic crisp sets and neutrosophic crisp continuity relations. Some related properties have been established with example. It 's hoped that our work will enhance this study in neutrosophic set theory.

References

[1] K. Atanassov, intuitionistic fuzzy sets, in V.Sgurev, ed., Vii ITKRS Session, Sofia(June 1983 central Sci. and Techn. Library, Bulg. Academy of Sciences (1984).
 [2] K. Atanassov, intuitionistic fuzzy sets, Fuzzy Sets and Systems 2087-96, (1986)
 [3] K. Atanassov, Review and new result on intuitionistic fuzzy sets, preprint IM-MFAIS-1-88, Sofia, (1988).
 [4] S. A. Alblowi, A.A. Salama and Mohmed Eisa, New Concepts of Neutrosophic Sets, International Journal of Mathematics and Computer Applications Research (IJMCAR), Vol. 4, Issue 1, (2014) 59-66.
 [5] I. M. Hanafy, A.A. Salama and K. Mahfouz, Correlation of Neutrosophic Data, International Refereed Journal of Engineering and Science (IRJES) , Vol.(1), Issue 2 .(2012), pp.39-33
 [6] I.M. Hanafy, A.A. Salama and K.M. Mahfouz,," Neutrosophic Classical Events and Its Probability" International Journal of Mathematics and Computer Applications Research(IJMCAR) Vol.(3),Issue 1,Mar (2013) ,pp.171-178.
 [7] A. A. Salama and S.A. Alblowi, "Generalized Neutrosophic Set and Generalized Neutrosophic

Spaces,"Journal Computer Sci. Engineering, Vol. (2) No. (7) (2012),pp.129-132 .
 [8] A. A. Salama and S. A. Alblowi, Neutrosophic Set and Neutrosophic Topological Spaces, ISOR J. Mathematics, Vol.(3), Issue(3), (2012), pp.31-35.
 [9] A.A. Salama, Neutrosophic Crisp Point & Neutrosophic Crisp Ideals, Neutrosophic Sets and Systems, Vol.1, No. 1, (2013) pp. 50-54.
 [10] A. A. Salama and F. Smarandache, Filters via Neutrosophic Crisp Sets, Neutrosophic Sets and Systems, Vol.1, No. 1, (2013), pp. 34-38.
 [11] A.A. Salama and S.A. Alblowi, Intuitionistic Fuzzy Ideals Spaces, Advances in Fuzzy Mathematics , Vol.(7), Number 1, (2012) pp. 51- 60.
 [12] A.A. Salama, and H.Elagamy, Neutrosophic Filters, International Journal of Computer Science Engineering and Information Technology Research (IJCSEITR), Vol.3, Issue(1),Mar 2013,(2013) pp 307-312.
 [13] A.A. Salama, F. Smarandache and V. Kroumov, Neutrosophic crisp Sets & Neutrosophic crisp Topological Spaces, Neutrosophic Sets and Systems, Vol.(2), pp25-30, (2014).
 [14] A.A. Salama, Mohamed Eisa and M. M. Abdelmoghny, Neutrosophic Relations Database, International Journal of Information Science and Intelligent System, 3(1) (2014).
 [15] A. A. Salama , Florentin Smarandache and S. A. Alblowi, New Neutrosophic Crisp Topological Concepts, Neutrosophic Sets and Systems, (Accepted), (2014) .
 [16] M. Bhowmik and M. Pal, Intuitionistic Neutrosophic Set Relations and Some of its Properties, Journal of Information and Computing Science, Vol.(5),No.3,(2010),pp.183-192.
 [17] F. Smarandache, Neutrosophy and Neutrosophic Logic, First International Conference on Neutrosophy , Neutrosophic Logic, Set, Probability, and Statistics University of New Mexico, Gallup, NM 87301, USA(2002).
 [18] F.Smarandache, A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic crisp Set, Neutrosophic Probability. American Research Press, Rehoboth, NM, (1999).
 [19] F. Smarandache, Neutrosophic set, a generalization of the intuitionistics fuzzy sets, Inter. J. Pure Appl. Math., 24 (2005), pp.287 – 297.
 [20] F. Smarandache, Introduction to neutrosophic measure, neutrosophic measure neutrosophic integral, and neutrosophic probability(2013). <http://fs.gallup.unm.edu/eBooks-otherformats.htm> EAN: 9781599732534
 [21] L.A. Zadeh, Fuzzy Sets, Inform and Control 8,(1965),pp.338-353.
 [22] A. Salama, S. Broumi and F. Smarandache, Neutrosophic Crisp Open Set and Neutrosophic Crisp Continuity via Neutrosophic Crisp Ideals, II. Information Engineering and Electronic Business, Vol.6, No.3,(2014) 1-8

Dr. Florentin Smarandache is a Professor of Mathematics at the University of New Mexico in USA. He published over 75

Some Types of Neutrosophic Crisp Sets and Neutrosophic Crisp Relations



books and 200 articles and notes in mathematics, physics, philosophy, psychology, rebus, literature. In mathematics his research is in number theory, non-Euclidean geometry, synthetic geometry, algebraic structures, statistics, neutrosophic logic and set (generalizations of fuzzy logic and set respectively), neutrosophic probability (generalization of classical and imprecise probability). Also, small contributions to nuclear and particle physics, information fusion, neutrosophy (a generalization of dialectics), law of sensations and stimuli, etc. He got the 2010 Telesio-Galilei Academy of Science Gold Medal, Adjunct Professor (equivalent to Doctor Honoris Causa) of Beijing Jiaotong University in 2011, and 2011 Romanian Academy Award for Technical Science (the highest in the country). Dr. W. B. Vasantha Kandasamy and Dr. Florentin Smarandache got the 2012 and 2011 New Mexico-Arizona Book Award for Algebraic Structures.



Dr. A.A. Salama (Ahmed Salama)
 Doctor and Lecturer in Mathematics and Computer Sciences Department in Faculty of Science in Port Said University. Associate Professor of Pure Mathematics & Computer Science in Baha College of Sciences, Saudi Arabia. Obtained Doctoral degree in 2001 in Pure Mathematics. 3. He published over 100 articles and notes in mathematics, computer science and Statistics.

- The first Arab to use the Neutrosophic concepts in these areas (computer Sci., Math, Statistics and Topology).
- A member of its Editorial Board to International Journal's Neutrosophic Set and Systems (USA).
- He published over 100 articles and notes in mathematics, computer science and Statistics.
- Reviewers of The Book MARIUS COMAN THE MATH ENCYCLOPEDIA OF SMARANDACHE TYPE NOTIONS . I. NUMBER THEORY Educational Publishing , 2013 by Marius Coman Education Publishing USA.
- A member of Editorial Board SMARANDACHE NOTIONS Journal's Vol.iii, ii, i, USA.
- Manager of the Quality Assurance Unit, Port Said Faculty of Science.
- Head of the Committee of Training and Community Service, Al-Baha Private College of Science.
- Educational Supervisor of Mathematics in the Zahraa Islamic for Language Schools, Mansoura for six years.
- Secretary-general of Topology Conference held in the Suez Canal University, 2007.
- Staff Member in the Higher Institute Tebah for Computer and Administrative Sciences, Maadi,

Cairo, Egypt.

- Head of the Board of Al-Haram Educational Periodical published in London.
- American Diploma Delegate, American Eagles Schools. 2005-2006.
- Main research points currently are Neutrosophic Mathematics, Computer Sciences and Statistics.



Said Broumi worked in Hassan II Casablanca university as an administrator. He worked in University for six years. He received his M. Sc in control system from Hassan II University Ain chok-Casablanca. A member of Editorial board to International Journal's Neutrosophic Set and Systems (USA). His research concentrates on soft set theory, fuzzy theory, intuitionistic fuzzy theory, neutrosophic theory, control systems. He has published 20 articles in international journals